

CSE 422 Lecture #13: Sampling and Estimation

This week, we will cover tools for making inferences based on random samples drawn from some distribution of interest (e.g., a distribution over voter priorities, customer behavior, ip addresses, etc.). Next lecture, we will see how to use sampling techniques to solve hard problems—both problems that inherently involve randomness, as well as those that do not.

As a warmup, to get into the probabilistic mindset, let's look at a cute and useful algorithm for drawing samples from a data stream.

1 Reservoir Sampling

Problem: How can one efficiently sample k uniformly random elements from a datastream of length $N \gg k$? The two main issues are 1) N might be huge—we certainly don't want to store a significant fraction of the stream in memory, 2) N might be unknown—we might not know the length of the stream ahead of time, but we still hope that whenever the stream ends, we have a uniformly random sample. (Recall that we are asking this problem in a similar “streaming” setting as that of the Count-Min-Sketch algorithm we saw earlier.)

As a motivating example, suppose a router might want to sample 1,0000 random IP addresses from a given day's traffic. The router obviously doesn't want to store all of the IP addresses, and the router doesn't know ahead of time the total amount of traffic that it will see on a given day.

Algorithm 1

RESERVOIR SAMPLING [VITTER '85]

Given a number k , and a datastream x_1, x_2, \dots of length greater than k :

- Put the first k elements of the stream into a “reservoir” $R = (x_1, \dots, x_k)$.
- For $i \geq k + 1$
 - With probability $\frac{k}{i}$ replace a random entry of R with x_i .
- At the end of the stream, return the reservoir R .

We now show that at any time $t \geq k$, the reservoir R consists of a uniformly random subset of k of the entries of x_1, \dots, x_t . To do this, let R_t denote the reservoir after the t th datastream element has been seen. It suffices to show that for all $t \geq i$, $\mathbb{P}[x_i \in R] = \frac{k}{t}$.

Claim 1.1 For all $t \geq i$, $\mathbb{P}[x_i \in R_t] = \frac{k}{t}$, where R_t denotes the reservoir after time t .

Proof: We prove this by induction on t . If $i \leq k$, the base case will be when $t = k$ and $\mathbb{P}[x_i \in R_k] = 1 = k/t$. If $i > k$, the base case will be $t = i$, and recall that the algorithm specifies that

we include x_i in R with probability exactly $k/i = k/t$. For the induction hypothesis, assume that the statement holds for some fixed value of $t \geq i$. We now consider $\mathbb{P}[x_i \in R_{t+1}]$ and leverage our induction hypothesis and conditional probability:

$$\begin{aligned} \mathbb{P}[x_i \in R_{t+1}] &= \mathbb{P}[x_i \in R_t] \cdot \mathbb{P}[x_i \text{ not replaced at time } t+1 | x_i \in R_t] \\ &= \frac{k}{t} \cdot \mathbb{P}[x_i \text{ not replaced at time } t+1 | x_i \in R_t] \quad \text{by our induction hyp.} \\ &= \frac{k}{t} \cdot \left(1 - \frac{k}{t+1} \cdot \frac{1}{k}\right) = \frac{k}{t+1}. \end{aligned}$$

Note that in the last line, we computed $\mathbb{P}[x_i \text{ not replaced at time } t+1 | x_i \in R_t]$ as $1 - \mathbb{P}[x_{t+1} \text{ replaces } x_i | x_i \in R_t] = 1 - \frac{k}{t+1} \frac{1}{k}$, where the $\frac{k}{t+1}$ term is the probability that x_{t+1} is stored in R_{t+1} , and the $1/k$ term is the probability that x_i is chosen as the random element of R_t to remove. ■

2 Basic Probability Tools: Our good friends Andrey [Markov] and Pafnuty [Chebyshev]

We begin with two of the most basic tools from probability theory that are extremely helpful for designing sampling schemes, and interpreting the significance of results gleaned from random samples.

Data is everywhere, and that there is a huge difference between what is possible with a careful, mathematically principled analysis, versus a sloppy/crude analysis. Just ask Nate Silver, who has essentially made a career out of properly analyzing publicly available political poll data (and, other public data, such as sports data, etc.). There is a link from the course page to a page on his website explaining his political prediction poll-aggregation system, in case you're interested.

Markov's inequality expresses the basic fact that

“at most 10% of the population can have an income that is more than $10 \times$ the average income of the population”:

Theorem 2.1 (Markov's Inequality) *For a real-valued random variable X satisfying $X \geq 0$, it holds that for any $c > 0$:*

$$\mathbb{P}[X \geq c\mathbb{E}[X]] \leq \frac{1}{c}.$$

Proof: Assume for the point of contradiction that $\mathbb{P}[X \geq c \cdot \mathbb{E}[X]] > \frac{1}{c}$, then $\mathbb{E}[X] > \frac{1}{c}c \cdot \mathbb{E}[X] = \mathbb{E}[X]$, which is a contradiction. ■

Markov's inequality tells us something very simple about a distribution over real numbers if all we know about the distribution is its expectation, and that it is non-negative. In many applications, it gives extremely weak bounds (e.g. the probability that a student's GPA is more than twice the average GPA is at most $1/2$ —not very surprising). Nevertheless, Markov's inequality is the key tool that lets us prove more powerful theorems that give sharper characterizations of distributions for which we have more information. For example, it lets us prove Chebyshev's inequality, which tells us that the probability that a random variable is more than c standard deviations from its expectation, is at most $1/c^2$.

Theorem 2.2 (Chebyshev's Inequality) For a real-valued random variable X , and any $c > 0$,

$$\mathbb{P} \left[|X - \mathbb{E}[X]| \geq c\sqrt{\mathbf{Var}[X]} \right] \leq \frac{1}{c^2}.$$

Proof: Chebyshev's inequality is proved by applying Markov's inequality to a cleverly chosen random variable. Let $Y = (X - \mathbb{E}[X])^2$. Note that Y is a perfectly good random variable, satisfying $Y \geq 0$. To apply Markov's inequality, we need to know $\mathbb{E}[Y]$. First, note that

$$\mathbb{E}[Y] = \mathbb{E} \left[(X - \mathbb{E}[X])^2 \right] = \mathbf{Var}[X].$$

Then we can calculate:

$$\begin{aligned} \mathbb{P} \left[|X - \mathbb{E}[X]| \geq c\sqrt{\mathbf{Var}[X]} \right] &= \mathbb{P}[Y \geq c^2 \mathbf{Var}[X]] \\ &= \mathbb{P}[Y \geq c^2 \mathbb{E}[Y]] \leq \frac{1}{c^2} \quad (\text{by Markov's inequality}) \end{aligned}$$

■

What is this good for? Lets see an example:

Example 2.3 How many people must we poll to estimate the percentage of people who support candidate C? Specifically, say we want our answer to be accurate to $\pm 1\%$, with probability at least $3/4$.

Let $p \in [0, 1]$ denote the true probability that a randomly chosen person supports candidate C. Suppose we poll n people, and output the fraction that support the candidate. Let X_1, \dots, X_n denote the 0/1 indicator random variables with X_i indicating that the i th person polled supports the candidate. Let $Z = \sum_i X_i$ denote the number of people polled who support the candidate.

$$\mathbf{Var}[Z] = \sum_i \mathbf{Var}[X_i] = n \cdot p(1 - p).$$

In the above calculation, we used 1) the fact that the variance of a sum of independent random variables is the sum of the variances, and 2) a quick calculation showing that if X is 1 with probability p , and 0 otherwise, $\mathbf{Var}[X] = p(1 - p)^2 + (1 - p)p^2 = p(1 - p)$.

An error in our estimate of 1% corresponds to estimating $\mathbb{E}[Z]$ up to an error of $0.01 \cdot n$. From Chebyshev's inequality,

$$\begin{aligned} \mathbb{P}[|Z - \mathbb{E}[Z]| \geq 0.01n] &= \mathbb{P} \left[|Z - \mathbb{E}[Z]| \geq \frac{0.01n}{\sqrt{\mathbf{Var}[Z]}} \sqrt{\mathbf{Var}[Z]} \right] \\ &= \mathbb{P} \left[|Z - \mathbb{E}[Z]| \geq \frac{0.01\sqrt{n}}{\sqrt{p(1 - p)}} \sqrt{\mathbf{Var}[Z]} \right] \\ &\leq \frac{p(1 - p)}{n/100^2} \quad \text{from Chebyshev's inequality.} \end{aligned}$$

For any $p \in [0, 1]$, the numerator $p(1 - p) \leq 1/4$, and hence if $n \geq 100^2$, this probability is at most $1/4$, as desired. Note that if we know that p is very small (i.e., $p < 0.1$) or very close to 1, then the numerator $p(1 - p)$ is very small, and the required number of samples is correspondingly smaller.

What we have just shown is that, in general, to estimate the expectation of a 0/1 random variable to error $\pm \epsilon$, one needs roughly $O(1/\epsilon^2)$ independent samples.

If Chebyshev’s inequality tells us something about a distribution given a bound on its variance, you might suspect that, given information on a distributions’ higher moments (i.e., $\mathbb{E}[X^3]$, $\mathbb{E}[X^4]$, etc.), one might be able to obtain even sharper insights into the distribution. This intuition is true, and can be made rigorous via the same approach as the proof of Chebyshev’s: apply Markov’s inequality to a cleverly chosen random variable, corresponding to one of the higher moments of the distribution you care about. This is also how one proves “central limit” style inverse-exponential bounds on tail probabilities (i.e., showing that the probability you flip fewer than 400 ‘heads’ in 1000 tosses of a fair coin is miniscule.)

3 Importance Sampling

Let us mention one setting where one can redesign a sampling protocol to improve the variance: *Importance sampling*. The basic idea is that we can estimate properties of distribution A based on samples from distribution B . Sometimes, A and B are fixed, and sometimes we can design distribution B to let us answer questions about A using fewer samples than we would need if we were to directly sample from A . This savings is achieved by having distribution B place higher weight on the “important” elements of the domain. The following example will clarify this high level intuition.

Example 3.1 Suppose we want to estimate the average income of the population. We know that the distribution of incomes has a long tail; i.e., there is a small fraction of people who have very large incomes, and hence have a large effect on the average income. This long, but important tail, is what makes the estimation task hard—if we don’t get any samples from the tail, then our estimate will likely be too low, but we need to take many samples in order to get any representatives from the tail.

We can use importance sampling as follows: Suppose we know that computer scientists have higher income than average, and that computer scientists compose exactly a 0.05 fraction of the total population. Rather than taking n random samples from the population, suppose we take $0.8n$ samples of non-computer scientists, and $0.2n$ samples of random computer scientists. If a_1 is the average salary of the non-computer scientists in our sample, and a_2 is the average salary of the sample of computer scientists, we can estimate the population average salary as:

$$\widehat{\text{avg}} = 0.95a_1 + 0.05a_2.$$

The benefit of this is that by taking more samples from higher earners, we obtained a better estimate of this important tail. Of course, the final calculation needs to reweight these two samples so that our answer is still an unbiased estimate of the overall population average.

Formally, how does one characterize the improvement that we obtained by over sampling the computer scientists? It does not change the expectation of our answer—our estimate is still the unbiased estimate of the population average. The benefit is that we have reduced the variance of our estimate, by focussing our samples on the important portion of the distribution (important with respect to the property that we care about—in this case, estimating the mean).

4 Knowing the unknowns: Estimating the missing mass

Thus far we have discussed interpreting random samples drawn from distributions over $0/1$, or distributions over real numbers (as in estimating average salary). In many applications, particularly in natural language processing, and genetics, one obtains samples from a distribution that is supported on a huge number of incomparable items. For example, a distribution over words, or a distribution over rare genetic variants. In these settings, one might not even know the support of the distribution.

While it is generally not possible to know what elements of the support are missing from a given sample, we can still hope to estimate the fraction of the distribution that is composed of the missing elements. One concrete question is the following:

Given a set of random samples drawn from a distribution, what is the probability that the next sample we draw is a “new” domain element that we have not seen previously? Equivalently, what is the amount of probability mass in the distribution composed of domain elements that are not represented in our sample (i.e. the “missing mass”)?

This question was first studied by I.J. Good and Alan Turing, in their work at Bletchley Park during WWII. (They wanted to estimate the probability that the next enigma machine ciphertext would be a previously unseen ciphertext.) They proposed the following extremely simple estimate of the missing mass, known as the *Good–Turing* frequency estimation scheme. Given n independent draws from a distribution, they proposed the following estimate:

$$\mathbb{P}[\text{next draw is something new}] \approx \frac{\# \text{ elmts seen exactly once}}{n}.$$

There is a large literature analyzing and extending this basic estimate—just search for “Good–Turing frequency estimation.”

Example 4.1 Assuming radio stations play songs in a random order, if you have been listening to the radio for the past 5 hours, the probability that the next song you hear is a song you haven’t heard in the past 5 hours can be estimated as the fraction of songs that you have heard exactly once in the past 5 hours.

Example 4.2 An estimate of the probability that the next word you read is a new word that you haven’t seen before is the number of words that you have seen exactly once, divided by the total number of words that you have seen.

Here is a quick sketch/derivation of this estimate: Assume that we have n samples from a distribution p supported on some unknown domain X , with $p(x)$ denoting the probability that the

distribution assigns to element $x \in X$.

$$\begin{aligned} \text{fraction unseen probability mass} &= \mathbb{P}[\text{next draw is new}] \\ &= \sum_{x \in X} p(x) \cdot \mathbb{P}[x \text{ unseen}] \\ &= \sum_{x \in X} p(x)(1 - p(x))^n \\ &= \frac{1}{n} \sum_{x \in X} n \cdot p(x)(1 - p(x))^n \\ &\approx \frac{1}{n} \sum_{x \in X} n \cdot p(x)(1 - p(x))^{n-1} \\ &= \frac{\#\mathbb{E}[\text{elmts seen exactly once}]}{n} \end{aligned}$$