


## 7.5 Bipartite Matching

---

# Matching

## Matching.


- Input: undirected graph  $G = (V, E)$ .
- $M \subseteq E$  is a **matching** if each node appears in at most edge in  $M$ .
- Max matching: find a max cardinality matching.



# Bipartite Matching

## Bipartite matching.

- Input: undirected, **bipartite** graph  $G = (L, R, E)$ .
- $M \subseteq E$  is a **matching** if each node appears in at most one edge in  $M$ .
- Max matching: find a max cardinality matching.




matching  
1-2', 3-1', 4-5'

# Bipartite Matching

## Bipartite matching.

- Input: undirected, **bipartite** graph  $G = (L, R, E)$ .
- $M \subseteq E$  is a **matching** if each node appears in at most one edge in  $M$ .
- Max matching: find a max cardinality matching.




max matching  
1-1', 2-2', 3-3' 4-4'

# Bipartite Matching

## Max flow formulation.

- Create digraph  $G' = (L, R, \{s, t\}, E')$ .
- Direct all edges from  $L$  to  $R$ , and assign infinite (or unit) capacity.
- Add source  $s$ , and unit capacity edges from  $s$  to each node in  $L$ .
- Add sink  $t$ , and unit capacity edges from each node in  $R$  to  $t$ .




# Bipartite Matching: Proof of Correctness

**Theorem.** Max cardinality matching in  $G$  = value of max flow in  $G'$ .

**Pf.**  $\leq$

- Given max matching  $M$  of cardinality  $k$ .
- Consider flow  $f$  that sends 1 unit along each of  $k$  paths.
- $f$  is a flow, and has cardinality  $k$ .





# Bipartite Matching: Proof of Correctness

**Theorem.** Max cardinality matching in  $G$  = value of max flow in  $G'$ .

**Pf.  $\geq$**

- Let  $f$  be a max flow in  $G'$  of value  $k$ .
- Integrality theorem  $\Rightarrow$   $k$  is integral and can assume  $f$  is 0-1.
- Consider  $M$  = set of edges from  $L$  to  $R$  with  $f(e) = 1$ .
  - each node in  $L$  and  $R$  participates in at most one edge in  $M$
  - $|M| = k$ : consider cut  $(L \cup s, R \cup t)$  .



# Perfect Matching

**Def.** A matching  $M \subseteq E$  is **perfect** if each node appears in exactly one edge in  $M$ .

**Q.** When does a bipartite graph have a perfect matching?

Structure of bipartite graphs with perfect matchings.


- Clearly we must have  $|L| = |R|$ .
- What other conditions are necessary?
- What conditions are sufficient?

# Perfect Matching

**Notation.** Let  $S$  be a subset of nodes, and let  $N(S)$  be the set of nodes adjacent to nodes in  $S$ .

**Observation.** If a bipartite graph  $G = (L, R, E)$ , has a perfect matching, then  $|N(S)| \geq |S|$  for all subsets  $S \subseteq L$ .

**Pf.** Each node in  $S$  has to be matched to a different node in  $N(S)$ .




No perfect matching:  
 $S = \{ 2, 4, 5 \}$   
 $N(S) = \{ 2', 5' \}.$

# Marriage Theorem

**Marriage Theorem.** [Frobenius 1917, Hall 1935] Let  $G = (L, R, E)$  be a bipartite graph with  $|L| = |R|$ . Then,  $G$  has a perfect matching iff  $|N(S)| \geq |S|$  for all subsets  $S \subseteq L$ .

**Pf.**  $\Rightarrow$  This was the previous observation.




No perfect matching:  
 $S = \{ 2, 4, 5 \}$   
 $N(S) = \{ 2', 5' \}.$

# Proof of Marriage Theorem

Pf.  $\Leftarrow$  Suppose  $G$  does not have a perfect matching.

- Formulate as a max flow problem and let  $(A, B)$  be min cut in  $G'$ .
- By max-flow min-cut,  $\text{cap}(A, B) < |L|$ .
- Define  $L_A = L \cap A$ ,  $L_B = L \cap B$ ,  $R_A = R \cap A$ .
- $\text{cap}(A, B) = |L_B| + |R_A|$ .
- Since min cut can't use  $\infty$  edges:  $N(L_A) \subseteq R_A$ .
- $|N(L_A)| \leq |R_A| = \text{cap}(A, B) - |L_B| < |L| - |L_B| = |L_A|$ .
- Choose  $S = L_A$ .



$L_A = \{2, 4, 5\}$   
 $L_B = \{1, 3\}$   
 $R_A = \{2', 5'\}$   
 $N(L_A) = \{2', 5'\}$

# Bipartite Matching: Running Time

Which max flow algorithm to use for bipartite matching?


- Generic augmenting path:  $O(m \text{ val}(f^*)) = O(mn)$ .
- Capacity scaling:  $O(m^2 \log C) = O(m^2)$ .
- Shortest augmenting path:  $O(m n^{1/2})$ .

Non-bipartite matching.

- Structure of non-bipartite graphs is more complicated, but well-understood. [Tutte-Berge, Edmonds-Galai]
- Blossom algorithm:  $O(n^4)$ . [Edmonds 1965]
- Best known:  $O(m n^{1/2})$ . [Micali-Vazirani 1980]


# Maximum Flow Problem

Max flow problem. Find s-t flow of maximum value.



# Minimum Cut Problem

Min s-t cut problem. Find an s-t cut of minimum capacity.



## 7.6 Disjoint Paths


---

# Edge Disjoint Paths

**Disjoint path problem.** Given a digraph  $G = (V, E)$  and two nodes  $s$  and  $t$ , find the max number of edge-disjoint  $s$ - $t$  paths.

**Def.** Two paths are **edge-disjoint** if they have no edge in common.

**Ex:** communication networks.




# Edge Disjoint Paths

**Disjoint path problem.** Given a digraph  $G = (V, E)$  and two nodes  $s$  and  $t$ , find the max number of edge-disjoint  $s$ - $t$  paths.


**Def.** Two paths are **edge-disjoint** if they have no edge in common.

**Ex:** communication networks.



# Edge Disjoint Paths

Max flow formulation: assign unit capacity to every edge.




**Theorem.** Max number edge-disjoint s-t paths equals max flow value.

Pf.  $\leq$

- Suppose there are  $k$  edge-disjoint paths  $P_1, \dots, P_k$ .
- Set  $f(e) = 1$  if  $e$  participates in some path  $P_i$ ; else set  $f(e) = 0$ .
- Since paths are edge-disjoint,  $f$  is a flow of value  $k$ .  $\cdot$

# Edge Disjoint Paths

Max flow formulation: assign unit capacity to every edge.



**Theorem.** Max number edge-disjoint s-t paths equals max flow value.

**Pf.**  $\geq$


- Suppose max flow value is  $k$ .
- Integrality theorem  $\Rightarrow$  there exists 0-1 flow  $f$  of value  $k$ .
- Consider edge  $(s, u)$  with  $f(s, u) = 1$ .
  - by conservation, there exists an edge  $(u, v)$  with  $f(u, v) = 1$
  - continue until reach  $t$ , always choosing a new edge
- Produces  $k$  (not necessarily simple) edge-disjoint paths. ▪

can eliminate cycles to get simple paths if desired

# Network Connectivity

**Network connectivity.** Given a digraph  $G = (V, E)$  and two nodes  $s$  and  $t$ , find min number of edges whose removal disconnects  $t$  from  $s$ .

**Def.** A set of edges  $F \subseteq E$  **disconnects  $t$  from  $s$**  if all  $s$ - $t$  paths uses at least on edge in  $F$ .





# Edge Disjoint Paths and Network Connectivity

**Theorem.** [Menger 1927] The max number of edge-disjoint  $s$ - $t$  paths is equal to the min number of edges whose removal disconnects  $t$  from  $s$ .

Pf.  $\leq$

- Suppose the removal of  $F \subseteq E$  disconnects  $t$  from  $s$ , and  $|F| = k$ .
- All  $s$ - $t$  paths use at least one edge of  $F$ . Hence, the number of edge-disjoint paths is at most  $k$ .





# Disjoint Paths and Network Connectivity

**Theorem.** [Menger 1927] The max number of edge-disjoint s-t paths is equal to the min number of edges whose removal disconnects t from s.

Pf.  $\geq$

- Suppose max number of edge-disjoint paths is  $k$ .
- Then max flow value is  $k$ .
- Max-flow min-cut  $\Rightarrow$  cut  $(A, B)$  of capacity  $k$ .
- Let  $F$  be set of edges going from  $A$  to  $B$ .
- $|F| = k$  and disconnects  $t$  from  $s$ . ▪



## 7.10 Image Segmentation

---

# Image Segmentation

## Image segmentation.


- Central problem in image processing.
- Divide image into coherent regions.

**Ex:** Three people standing in front of complex background scene.  
Identify each person as a coherent object.

# Image Segmentation

## Foreground / background segmentation.

- Label each pixel in picture as belonging to foreground or background.
- $V$  = set of pixels,  $E$  = pairs of neighboring pixels.
- $a_i > 0$  is likelihood pixel  $i$  in foreground.
- $b_i > 0$  is likelihood pixel  $i$  in background.
- $p_{ij} > 0$  is separation penalty for labeling one of  $i$  and  $j$  as foreground, and the other as background.



## Goals.

- Accuracy: if  $a_i > b_i$  in isolation, prefer to label  $i$  in foreground.
- Smoothness: if many neighbors of  $i$  are labeled foreground, we should be inclined to label  $i$  as foreground.

- Find partition  $(A, B)$  that maximizes:
 
$$\sum_{i \in A} a_i + \sum_{j \in B} b_j - \sum_{\substack{(i,j) \in E \\ |A \cap \{i,j\}| = 1}} p_{ij}$$

$\nearrow$   
 foreground

$\nwarrow$   
 background

# Image Segmentation

Formulate as min cut problem.

- Maximization.
- No source or sink.
- Undirected graph.

Turn into minimization problem.

- Maximizing 
$$\sum_{i \in A} a_i + \sum_{j \in B} b_j - \sum_{\substack{(i,j) \in E \\ |A \cap \{i,j\}| = 1}} p_{ij}$$

is equivalent to maximizing



$$\underbrace{\left( \sum_{i \in V} a_i + \sum_{j \in V} b_j \right)}_{\text{a constant}} - \sum_{i \in B} a_i - \sum_{j \in A} b_j - \sum_{\substack{(i,j) \in E \\ |A \cap \{i,j\}| = 1}} p_{ij}$$

- or alternatively minimizing 
$$\sum_{j \in B} a_j + \sum_{i \in A} b_i + \sum_{\substack{(i,j) \in E \\ |A \cap \{i,j\}| = 1}} p_{ij}$$

# Image Segmentation

Formulate as min cut problem.

- $G' = (V', E')$ .
- Add source to correspond to foreground;  
add sink to correspond to background
- Use two anti-parallel edges instead of undirected edge.



# Image Segmentation


Consider min cut  $(A, B)$  in  $G'$ .

- $A$  = foreground.

$$cap(A, B) = \sum_{j \in B} a_j + \sum_{i \in A} b_i + \sum_{\substack{(i,j) \in E \\ i \in A, j \in B}} p_{ij}$$

← if  $i$  and  $j$  on different sides,  $p_{ij}$  counted exactly once

- Precisely the quantity we want to minimize.



## 7.11 Project Selection

---

# Project Selection

can be positive or negative



## Projects with prerequisites.


- Set  $P$  of possible projects. Project  $v$  has associated revenue  $p_v$ .
  - some projects generate money: create interactive e-commerce interface, redesign web page
  - others cost money: upgrade computers, get site license
- Set of prerequisites  $E$ . If  $(v, w) \in E$ , can't do project  $v$  and unless also do project  $w$ .
- A subset of projects  $A \subseteq P$  is **feasible** if the prerequisite of every project in  $A$  also belongs to  $A$ .

**Project selection.** Choose a feasible subset of projects to maximize revenue.


# Project Selection: Prerequisite Graph

## Prerequisite graph.

- Include an edge from  $v$  to  $w$  if can't do  $v$  without also doing  $w$ .
- $\{v, w, x\}$  is feasible subset of projects.
- $\{v, x\}$  is infeasible subset of projects.



feasible




infeasible

# Project Selection: Min Cut Formulation

## Min cut formulation.

- Assign capacity to all prerequisite edge.
- Add edge  $(s, v)$  with capacity  $p_v$  if  $p_v > 0$ .
- Add edge  $(v, t)$  with capacity  $-p_v$  if  $p_v < 0$ .
- For notational convenience, define  $p_s = p_t = 0$ .




# Project Selection: Min Cut Formulation

**Claim.**  $(A, B)$  is min cut iff  $A - \{s\}$  is optimal set of projects.

- Infinite capacity edges ensure  $A - \{s\}$  is feasible.


- Max revenue because:
 
$$\begin{aligned} \text{cap}(A, B) &= \sum_{v \in B: p_v > 0} p_v + \sum_{v \in A: p_v < 0} (-p_v) \\ &= \underbrace{\sum_{v: p_v > 0} p_v}_{\text{constant}} - \sum_{v \in A} p_v \end{aligned}$$



# Open Pit Mining

Open-pit mining. (studied since early 1960s)

- Blocks of earth are extracted from surface to retrieve ore.
- Each block  $v$  has net value  $p_v = \text{value of ore} - \text{processing cost}$ .
- Can't remove block  $v$  before  $w$  or  $x$ .




## 7.12 Baseball Elimination

"See that thing in the paper last week about Einstein? . . . Some reporter asked him to figure out the mathematics of the pennant race. You know, one team wins so many of their remaining games, the other teams win this number or that number. What are the myriad possibilities? Who's got the edge?"

"The hell does he know?"

"Apparently not much. He picked the Dodgers to eliminate the Giants last Friday."

- *Don DeLillo, Underworld*



# Baseball Elimination

Team $i$	Wins $w_i$	Losses $l_i$	To play $r_i$	Against = $r_{ij}$			
				Atl	Phi	NY	Mon
Atlanta	83	71	8	-	1	6	1
Philly	80	79	3	1	-	0	2
New York	78	78	6	6	0	-	0
Montreal	77	82	3	1	2	0	-

Which teams have a chance of finishing the season with most wins?

- Montreal eliminated since it can finish with at most 80 wins, but Atlanta already has 83.
- $w_i + r_i < w_j \Rightarrow$  team  $i$  eliminated.
- Only reason sports writers appear to be aware of.
- Sufficient, but not necessary!

# Baseball Elimination

Team $i$	Wins $w_i$	Losses $l_i$	To play $r_i$	Against = $r_{ij}$			
				Atl	Phi	NY	Mon
Atlanta	83	71	8	-	1	6	1
Philly	80	79	3	1	-	0	2
New York	78	78	6	6	0	-	0
Montreal	77	82	3	1	2	0	-

Which teams have a chance of finishing the season with most wins?

- Philly can win 83, but still eliminated . . .
- If Atlanta loses a game, then some other team wins one.

**Remark.** Answer depends not just on **how many** games already won and left to play, but also on **whom** they're against.

# Baseball Elimination

TUESDAY, SEPTEMBER 10, 1996

San Francisco Chronicle

The Gate

Sports Online

► <http://www.sfgate.com>

## SPORTING G

# 49ers, Young Get Big Break



## Quarterback m

By Gary Swan  
Chronicle Staff Writer

The bye week has come at a perfect time for the 49ers and quarterback Steve Young. If they had a game next Sunday, there's a good chance Young would not play.

Put the pulled groin muscle on his up-

# Giants Officially Leave the NL West Race

By Nancy Gay  
Chronicle Staff Writer

With the smack of another National League West bat 500 miles away, the Giants' run at the division title ended last night, just as they were handing the visiting St. Louis Cardinals an even bigger lead in the NL Central.

CARDINALS 6  
GIANTS 2

In San Diego, Greg Vaughn's three-run homer in the eighth pushed the Padres over the Pirates and officially shoved the rest of the Giants' season into the back-ground. On the heels of their tedi-ous 6-2 loss before an announced crowd of 10,307 at Candlestick Park, the Giants fell 19½ games off the lead.

As it is, the worst the Padres' (80-65) can finish is 80-82. The Giants have fallen to 59-83 with 20

Financing in Place  
For Giants' New Stadium  
SEE PAGE B1, MAIN NEWS

games left; they cannot win 80 games. Coming off a miserable 2-8 mark on a three-city road trip that saw their road record drop to 27-47, the Giants were hoping to get off on the right foot in their long-est homestand of the year (15 games, 14 days).

"Where we are, you're going to be eliminated sooner or later," Baker said quietly. "But it doesn't alter the fact that we've still got to play ball. You've still got to play hard, the fans come out to watch you play. You've got to play for the fact of loving to play, no matter where you are in the standings.

"You've got to play the role of spoiler, to not make it easier on

GIANTS: Page D5 Col. 3

# Baseball Elimination


Baseball elimination problem.

- Set of teams  $S$ .
- Distinguished team  $s \in S$ .
- Team  $x$  has won  $w_x$  games already.
- Teams  $x$  and  $y$  play each other  $r_{xy}$  additional times.
- Is there any outcome of the remaining games in which team  $s$  finishes with the most (or tied for the most) wins?

# Baseball Elimination: Max Flow Formulation

Can team 3 finish with most wins?


- Assume team 3 wins all remaining games  $\Rightarrow w_3 + r_3$  wins.
- Divvy remaining games so that all teams have  $< w_3 + r_3$  wins.



# Baseball Elimination: Max Flow Formulation

**Theorem.** Team 3 is not eliminated iff max flow saturates all edges leaving source.

- Integrality theorem  $\Rightarrow$  each remaining game between  $x$  and  $y$  added to number of wins for team  $x$  or team  $y$ .
- Capacity on  $(x, t)$  edges ensure no team wins too many games.



# Baseball Elimination: Explanation for Sports Writers

Team $i$	Wins $w_i$	Losses $l_i$	To play $r_i$	Against = $r_{ij}$				
				NY	Bal	Bos	Tor	Det
NY	75	59	28	-	3	8	7	3
Baltimore	71	63	28	3	-	2	7	4
Boston	69	66	27	8	2	-	0	0
Toronto	63	72	27	7	7	0	-	-
Detroit	49	86	27	3	4	0	0	-

AL East: August 30, 1996

Which teams have a chance of finishing the season with most wins?

- Detroit could finish season with  $49 + 27 = 76$  wins.

# Baseball Elimination: Explanation for Sports Writers

Team $i$	Wins $w_i$	Losses $l_i$	To play $r_i$	Against = $r_{ij}$				
				NY	Bal	Bos	Tor	Det
NY	75	59	28	-	3	8	7	3
Baltimore	71	63	28	3	-	2	7	4
Boston	69	66	27	8	2	-	0	0
Toronto	63	72	27	7	7	0	-	-
Detroit	49	86	27	3	4	0	0	-

AL East: August 30, 1996

Which teams have a chance of finishing the season with most wins?

- Detroit could finish season with  $49 + 27 = 76$  wins.

Certificate of elimination.  $R = \{NY, Bal, Bos, Tor\}$

- Have already won  $w(R) = 278$  games.
- Must win at least  $r(R) = 27$  more.
- Average team in  $R$  wins at least  $305/4 > 76$  games.

# Baseball Elimination: Explanation for Sports Writers

Certificate of elimination.

$$T \subseteq S, \quad w(T) := \overbrace{\sum_{i \in T} w_i}^{\# \text{ wins}}, \quad g(T) := \overbrace{\sum_{\{x,y\} \subseteq T} g_{x,y}}^{\# \text{ remaining games}},$$

LB on avg # games won  
If  $\frac{w(T) + g(T)}{|T|} > w_z + g_z$  then  $z$  is **eliminated** (by subset  $T$ ).


**Theorem.** [Hoffman-Rivlin 1967] Team  $z$  is eliminated iff there exists a subset  $T^*$  that eliminates  $z$ .

**Proof idea.** Let  $T^*$  = team nodes on source side of min cut.

# Baseball Elimination: Explanation for Sports Writers

## Pf of theorem.

- Use max flow formulation, and consider min cut  $(A, B)$ .
- Define  $T^*$  = team nodes on source side of min cut.
- Observe  $x-y \in A$  iff both  $x \in T^*$  and  $y \in T^*$ .
  - infinite capacity edges ensure if  $x-y \in A$  then  $x \in A$  and  $y \in A$
  - if  $x \in A$  and  $y \in A$  but  $x-y \in B$ , then adding  $x-y$  to  $A$  decreases capacity of cut



# Baseball Elimination: Explanation for Sports Writers

Pf of theorem.

- Use max flow formulation, and consider min cut  $(A, B)$ .
- Define  $T^*$  = team nodes on source side of min cut.
- Observe  $x-y \in A$  iff both  $x \in T^*$  and  $y \in T^*$ .
- $g(S - \{z\}) > \text{cap}(A, B)$

$$\begin{aligned} & \text{capacity of game edges leaving } s && \text{capacity of team edges leaving } s \\ = & \overbrace{g(S - \{z\}) - g(T^*)} && + \overbrace{\sum_{x \in T^*} (w_z + g_z - w_x)} \\ = & g(S - \{z\}) - g(T^*) && - w(T^*) + |T^*|(w_z + g_z) \end{aligned}$$

- Rearranging terms:  $w_z + g_z < \frac{w(T^*) + g(T^*)}{|T^*|}$  .