

degrees

$\deg(v)$ = number of vertices adjacent to v

degrees

deg(v) = number of vertices adjacent to v

Claim: If a graph has m edges, then

$$\sum_v \deg(v) = 2m.$$

Proof: Every edge $\{u,v\}$ contributes exactly 2 to the left hand side, 1 to $\deg(u)$, and 1 to $\deg(v)$.

degrees

Claim: In a party, the number of people who shake hands with an odd number of people must be even.


Proof: Construct a graph. Each vertex represents person, put edge between 2 people if they shake hands.

Need to prove: #vertices with odd degree is even.

$$2m = \sum \deg(v) = \underbrace{\sum \deg(v)}_{\text{even}} + \underbrace{\sum \deg(v)}_{\text{even}}$$

Thus $\sum \deg(v)$ is even, so # vertices with odd degree is even.

Claim: If a graph has no cycles, it must have a vertex v of degree ≤ 1 .



Path: A sequence of distinct vertices where each vertex is connected to the next by an edge.

Cycle: A path of length > 1 such that the first vertex is connected to the last one.

1,2,5,3 is a cycle

6,4,2,1,3,5,2,4 is not a cycle


Claim: If a graph has no cycles, it must have a vertex v of degree ≤ 1 .

Proof:

Suppose not.

Then there is a graph G that has no cycles, and yet every vertex in the graph has degree > 1 .

There must be v_0, v_1, \dots, v_n , a sequence of $n+1$ vertices such that $v_{i-1} \neq v_{i+1}$, for all i , and adjacent vertices have an edge.



Intuition: this should not be possible! Let's use the degrees of the vertices to find a cycle.

Note: The argument should work for every such graph!

Claim: If a graph has no cycles, it must have a vertex v of degree ≤ 1 .

Proof:


Suppose not.

Then there is a graph G that has no cycles, and yet every vertex in the graph has degree > 1 .

There must be v_0, v_1, \dots, v_n , a sequence of $n+1$ vertices such that $v_{i-1} \neq v_{i+1}$, for all i , and adjacent vertices have an edge.

By the pigeonhole principle, there must $i < j$ s.t. $v_i = v_j$.

Let $i < j$ be the closest such pair. Then $v_i, v_{i+1}, \dots, v_{j-1}$ form a cycle.



Intuition: this should not be possible! Let's use the degrees of the vertices to find a cycle.

Note: The argument should work for every such graph!

Lemma: Every tree on n vertices has exactly $n-1$ edges.

Tree: A connected graph that has no cycles.


Proof: By induction on n .

Base case: $n=1$. Every graph with 1 vertex has $1-1 = 0$ edges. The claim holds.

Inductive step: $n > 1$. Let T be a tree with n vertices.

Since T has no cycles, there is a vertex v , $\deg(v) \leq 1$ by previous claim. $\deg(v)=1$ since T is connected means there are no degree 0 vertices.

$T' = T - v$ is connected and has no cycles, so it is a tree. T' has $n-1$ vertices. So by induction, T' has $n-2$ edges. Thus T has $n-1$ edges.



Mantel's Theorem: If a graph on $2n$ vertices has n^2+1 edges, then it must have a triangle (3-cycle).

Proof: By induction on n .

Mantel's Theorem: If a graph on $2n$ vertices has n^2+1 edges, then it must have a triangle (3-cycle).

Proof: By induction on n .

Base case: $n=1$. Every graph with 2 vertices has at most 1 edge, so it can never have $n^2+1 = 2$ edges. The claim holds vacuously.


Mantel's Theorem: If a graph on $2n$ vertices has n^2+1 edges, then it must have a triangle (3-cycle).

Proof: By induction on n .

Base case: $n=1$. Every graph with 2 vertices has at most 1 edge, so it can never have $n^2+1 = 2$ edges. The claim holds vacuously.

Inductive step: $n>1$.

Let $\{x,y\}$ be an edge of the graph, and let G' denote the subgraph using the rest of the vertices.



Mantel's Theorem: If a graph on $2n$ vertices has n^2+1 edges, then it must have a triangle (3-cycle).

Proof: By induction on n .


Base case: $n=1$. Every graph with 2 vertices has at most 1 edge, so it can never have $n^2+1 = 2$ edges. The claim holds vacuously.

Inductive step: $n>1$.

Let $\{x,y\}$ be an edge of the graph, and let G' denote the subgraph using the rest of the vertices.

Case 1: At least $(n-1)^2+1$ edges are in G' .

Then by the induction hypothesis, G' has a triangle.



Mantel's Theorem: If a graph on $2n$ vertices has n^2+1 edges, then it must have a triangle (3-cycle).

Proof: By induction on n .

Base case: $n=1$. Every graph with 2 vertices has at most 1 edge, so it can never have $n^2+1 = 2$ edges. The claim holds vacuously.

Inductive step: $n>1$.

Let $\{x,y\}$ be an edge of the graph, and let G' denote the subgraph using the rest of the vertices.

Case 1: At least $(n-1)^2+1$ edges are in G' .
Then by the induction hypothesis, G' has a triangle.

Case 2: At most $(n-1)^2$ edges are in G' .
There is one edge between x,y , so #edges from $\{x,y\}$ to $G' \geq n^2-(n-1)^2 = 2(n-1)+1$. But G' has $2(n-1)$ vertices, so there is vertex z such that $\{x,z\}, \{y,z\}$ are edges. Then x,y,z is a triangle.

