

Embedding the diamond graph in L_p and dimension reduction in L_1

James R. Lee* Assaf Naor
U.C. Berkeley and Microsoft Research Microsoft Research

Abstract

We show that any embedding of the level k diamond graph of Newman and Rabinovich [6] into L_p , $1 < p \leq 2$, requires distortion at least $\sqrt{k(p-1)+1}$. An immediate corollary is that there exist arbitrarily large n -point sets $X \subseteq L_1$ such that any D -embedding of X into ℓ_1^d requires $d \geq n^{\Omega(1/D^2)}$. This gives a simple proof of a recent result of Brinkman and Charikar [2] which settles the long standing question of whether there is an L_1 analogue of the Johnson-Lindenstrauss dimension reduction lemma [4].

1 The diamond graphs, distortion, and dimension

We recall the definition of the diamond graphs $\{G_k\}_{k=0}^\infty$ whose shortest path metrics are known to be uniformly bi-lipschitz equivalent to a subset of L_1 (see [3] for the L_1 embeddability of general series-parallel graphs). The diamond graphs were used in [6] to obtain lower bounds for the Euclidean distortion of planar graphs and similar arguments were previously used in a different context by Laakso [5].

G_0 consists of a single edge of length 1. G_i is obtained from G_{i-1} as follows. Given an edge $(u, v) \in E(G_{i-1})$, it is replaced by a quadrilateral u, a, v, b with edge lengths 2^{-i} . In what follows, (u, v) is called an edge of level $i-1$, and (a, b) is called the level i anti-edge corresponding to (u, v) . Our main result is a lower bound on the distortion necessary to embed G_k into L_p , for $1 < p \leq 2$.

Theorem 1.1. *For every $1 < p \leq 2$, any embedding of G_k into L_p incurs distortion at least $\sqrt{1+(p-1)k}$.*

The following corollary shows that the diamond graphs cannot be well-embedded into low-dimensional ℓ_1 spaces. In particular, an L_1 analogue of the Johnson-Lindenstrauss dimension reduction lemma does not exist. The same graphs were used in [2] as an example which shows the impossibility of dimension reduction in L_1 . Our proof is different and, unlike the linear programming based argument appearing there, relies on geometric intuition. We proceed by observing that a lower bound on the rate of decay of the distortion as $p \rightarrow 1$ yields a lower bound on the required dimension in ℓ_1 .

Corollary 1.2. *For every $n \in \mathbb{N}$, there exists an n -point subset $X \subseteq L_1$ such that for every $D > 1$, if X D -embeds into ℓ_1^d , then $d \geq n^{\Omega(1/D^2)}$.*

*Work partially supported by NSF grant CCR-0121555 and an NSF Graduate Research Fellowship.

Proof. Since ℓ_1^d is $O(1)$ -isomorphic to ℓ_p^d when $p = 1 + \frac{1}{\log d}$ and G_k is $O(1)$ -equivalent to a subset $X \subseteq L_1$, it follows that $\sqrt{1 + \frac{k}{\log d}} = O(D)$. Noting that $k = \Omega(\log n)$ completes the proof. \square

2 Proof

The proof is based on the following inequality. The case $p = 2$ is the well known “short diagonals lemma” which was central to the argument in [5, 6].

Lemma 2.1. *Fix $1 < p \leq 2$ and $x, y, z, w \in L_p$. Then,*

$$\|y - z\|_p^2 + (p - 1)\|x - w\|_p^2 \leq \|x - y\|_p^2 + \|y - w\|_p^2 + \|w - z\|_p^2 + \|z - x\|_p^2.$$

Proof. For every $a, b \in L_p$, $\|a + b\|_p^2 + (p - 1)\|a - b\|_p^2 \leq 2(\|a\|_p^2 + \|b\|_p^2)$. A simple proof of this classical fact can be found, for example, in [1]. Now,

$$\|y - z\|_p^2 + (p - 1)\|y - 2x + z\|_p^2 \leq 2\|y - x\|_p^2 + 2\|x - z\|_p^2$$

and

$$\|y - z\|_p^2 + (p - 1)\|y - 2w + z\|_p^2 \leq 2\|y - w\|_p^2 + 2\|w - z\|_p^2.$$

Averaging these two inequalities yields

$$\|y - z\|_p^2 + (p - 1) \frac{\|y - 2x + z\|_p^2 + \|y - 2w + z\|_p^2}{2} \leq \|x - y\|_p^2 + \|y - w\|_p^2 + \|w - z\|_p^2 + \|z - x\|_p^2.$$

The required inequality follows by convexity. \square

Lemma 2.2. *Let A_i denote the set of anti-edges at level i and let $\{s, t\} = V(G_0)$, then for $1 < p \leq 2$ and any $f : G_k \rightarrow L_p$,*

$$\|f(s) - f(t)\|_p^2 + (p - 1) \sum_{i=1}^k \sum_{(x,y) \in A_i} \|f(x) - f(y)\|_p^2 \leq \sum_{(x,y) \in E(G_k)} \|f(x) - f(y)\|_p^2.$$

Proof. Let (a, b) be an edge of level i and (c, d) its corresponding anti-edge. By Lemma 2.1, $\|f(a) - f(b)\|_p^2 + (p - 1)\|f(c) - f(d)\|_p^2 \leq \|f(a) - f(c)\|_p^2 + \|f(b) - f(c)\|_p^2 + \|f(d) - f(a)\|_p^2 + \|f(d) - f(b)\|_p^2$. Summing over all such edges and all $i = 0, \dots, k - 1$ yields the desired result by noting that the terms $\|f(x) - f(y)\|_p^2$ corresponding to $(x, y) \in E(G_i)$ cancel for $i = 1, \dots, k - 1$. \square

The main theorem now follows easily.

Proof of Theorem 1.1. Let $f : G_k \rightarrow L_p$ be a non-expansive D -embedding. Since $|A_i| = 4^{i-1}$ and the length of a level i anti-edge is 2^{1-i} , applying Lemma 2.2 yields $\frac{1+(p-1)k}{D^2} \leq 1$. \square

References

- [1] K. Ball, E. A. Carlen and E. Lieb. *Sharp uniform convexity and smoothness inequalities for trace norms*. Invent. Math. 115, 463-482 (1994).
- [2] B. Brinkman and M. Charikar. *On the Impossibility of Dimension Reduction in ℓ_1* . In Proceedings of the 44th Annual IEEE Conference on Foundations of Computer Science. ACM, 2003.
- [3] A. Gupta, I. Newman, Y. Rabinovich and A. Sinclair. *Cuts, trees and ℓ_1 embeddings*. In Proceedings of the 40th Annual Symposium on Foundations of Computer Science. ACM, 1999.
- [4] W. B. Johnson and J. Lindenstrauss. *Extensions of Lipschitz mappings into a Hilbert space*. In *Conference in modern analysis and probability (New Haven, Conn., 1982)*, pages 189-206. Amer. Math. Soc., Providence, RI, 1984.
- [5] T. J. Laakso. *Ahlfors Q -regular spaces with arbitrary $Q > 1$ admitting weak Poincaré inequality*. Geom. Funct. Anal., 10(1):111-123, 2000.
- [6] I. Newman and Y. Rabinovich. *A Lower Bound on the Distortion of Embedding Planar Metrics into Euclidean Space*. Discrete Computational Geometry, 29 no. 1, 77-81 (2003).