

The intrinsic dimensionality of graphs

Robert Krauthgamer*

James R. Lee†

June 16, 2006

Abstract

We resolve the following conjecture raised by Levin together with Linial, London, and Rabinovich [Combinatorica, 1995]. For a graph G , let $\dim(G)$ be the smallest d such that G occurs as a (not necessarily induced) subgraph of \mathbb{Z}_∞^d , the infinite graph with vertex set \mathbb{Z}^d and an edge (u, v) whenever $\|u - v\|_\infty = 1$. The *growth rate* of G , denoted ρ_G , is the minimum ρ such that every ball of radius $r > 1$ in G contains at most r^ρ vertices. By simple volume arguments, $\dim(G) = \Omega(\rho_G)$. Levin conjectured that this lower bound is tight, i.e., that $\dim(G) = O(\rho_G)$ for every graph G .

Previously, it was unknown whether $\dim(G)$ could be bounded above by any function of ρ_G . We show that a weaker form of Levin's conjecture holds by proving that $\dim(G) = O(\rho_G \log \rho_G)$ for any graph G . We disprove, however, the specific bound of the conjecture and show that our upper bound is tight by exhibiting graphs for which $\dim(G) = \Omega(\rho_G \log \rho_G)$. For several special families of graphs (e.g., planar graphs), we salvage the strong form, showing that $\dim(G) = O(\rho_G)$. Our results extend to a variant of the conjecture for finite-dimensional Euclidean spaces posed by Linial and independently by Benjamini and Schramm.

1 Introduction

The geometry of graphs, a fascinating area of combinatorics concerned with the geometric representation of graphs, has found many algorithmic applications in recent years. A very fruitful and actively studied line of research involves embedding the metric of a weighted graph into some finite-dimensional real-normed space (see, for instance, the surveys [13] and [19, ch. 15]). In their seminal paper [16], Linial, London, and Rabinovich were the first to fully realize the algorithmic importance of low-distortion metric embeddings. However, their initial motivation was to understand the relationship between the *dimensionality* of a graph and its combinatorial properties.

The notion of dimensionality for a graph is usually based on a particular way of embedding the graph into some space that possesses an intrinsic dimension (e.g., a finite-dimensional Euclidean space). One then defines the dimension of a graph to be the least dimension into which it can be embedded. Several such notions have been extensively studied; see [18]. In [16], the authors wished to express the concept that graphs of “everywhere large diameter” should have low dimensionality. With the help of Leonid Levin, this concept was formalized as follows.

*IBM Almaden Research Center, 650 Harry Road, San Jose CA 95120, USA. Email: robi@almaden.ibm.com

†Computer Science Division, University of California, Berkeley, CA 94720. Supported by NSF grant CCR-0121555 and by an NSF Graduate Research Fellowship. Email: jrl@cs.berkeley.edu

Let \mathbb{Z}_∞^d be the infinite graph with vertex set \mathbb{Z}^d and an edge (u, v) for two vertices u and v whenever $\|u - v\|_\infty = 1$. For a graph $G = (V, E)$, define $\dim(G)$ to be the smallest d such that G occurs as a (not necessarily induced) subgraph of \mathbb{Z}_∞^d .

For a pair of vertices $u, v \in V$, we define $d_G(u, v)$ to be the distance between u and v in the shortest path metric on G . We denote by

$$B(v, r) = \{u \in V : d_G(u, v) \leq r\}$$

the closed ball of radius r in G centered at v , and define the *growth rate* of G to be

$$\rho_G = \inf\{\rho : |B(v, r)| \leq r^\rho \text{ for all } v \in V \text{ and } r > 1\}.$$

Equivalently, $\rho_G = \sup\{\frac{\log |B(v, r)|}{\log r} : v \in V, r > 1\}$. Notice that $\rho_{\mathbb{Z}_\infty^d} = \Theta(d)$, so by a simple counting argument, we must have $\dim(G) = \Omega(\rho_G)$. Levin, together with Linial, London, and Rabinovich [16], conjectured that $O(\rho_G)$ dimensions suffice.

Conjecture 1. *For any graph G with growth rate ρ_G , G occurs as a (not necessarily induced) subgraph of $\mathbb{Z}_\infty^{O(\rho_G)}$. In other words, $\dim(G) = \Theta(\rho_G)$.*

In [16], it was shown that Conjecture 1 holds for the k -dimensional hypercube and the complete binary tree, but nothing beyond these two special cases was known. Indeed, it was not known whether $\dim(G)$ could be upper bounded by *any* function of ρ_G , even in the seemingly simpler case when the graph is a tree. Linial [15] asked about a Euclidean analogue to this notion of dimensionality.

Question 2. *For a graph $G = (V, E)$, what is the minimum dimension d , denoted $\dim_2(G)$, such that there exists a mapping $\gamma : V \rightarrow \mathbb{R}^d$ with the following properties?*

1. $\|\gamma(u) - \gamma(v)\|_2 \geq 1$ for all $u \neq v \in V$, and
2. $\|\gamma(u) - \gamma(v)\|_2 \leq 2$ for all $(u, v) \in E$.

Itai Benjamini and Oded Schramm [personal communication, 2003] independently asked a similar question: Is it the case that, for every infinite graph G with $\rho(G) < \infty$, we have $\dim_2(G) < \infty$? In what follows, we resolve all these questions and give tight quantitative bounds. Linial remarked that the condition $\|\gamma(u) - \gamma(v)\|_2 \leq 2$ is somewhat arbitrary; indeed, we will see that it can be replaced by $\|\gamma(u) - \gamma(v)\|_2 \leq C$ for any fixed $C > 1$ while affecting the value of $\dim_2(G)$ by only a constant factor (that depends on C), see Section 6.

1.1 Results and techniques

In Section 2, we provide a self-contained proof of Levin’s conjecture for trees. We first perform a recursive decomposition of the tree. Each “level” of the decomposition is responsible for pairs of vertices whose distance (in the tree) falls into a certain range of the form $[r, r^2]$. For any single level, we use the probabilistic method to construct an embedding that handles all the respective pairs of vertices. We can embed each level separately and concatenate the $O(\log \log |V|)$ resulting mappings, but the dimension of the host lattice becomes too large. The key technique, which completes the proof, is a method of handling all the levels using the same coordinates. The solution involves “conservation” of randomness between levels—our hierarchical decomposition of the tree induces a partition of a corresponding probability space, and the probabilistic construction

at later levels is restricted to randomizing over only the remaining “untapped” randomness (an idea that reappears throughout the paper). This result extends to graphs whose induced simple cycles are of bounded length by known low-distortion embeddings of such graphs into trees, due to [5, 6].

In Section 3, we give a lower bound on the dimension necessary to embed low-degree expander graphs which shows that the strong form of Levin’s conjecture does not hold in general. In particular, we show that for a $\log n$ -degree expander G , $\dim(G) = \Omega(\rho_G \log \rho_G)$.

In Section 4, using different techniques, but many ideas from our proof for trees, we give a general upper bound on $\dim(G)$ in terms of certain graph decompositions. In this setting, choosing a good embedding for a level with high probability is more difficult; the approach we use is inspired by a technique of [21] (which was used there to embed planar graphs into Euclidean space with low distortion). Again, we must discover a delicate way of handling all the levels simultaneously. In Section 4.4, we employ the decomposition of [14] (see also [10]), combined with the results of Section 4, to prove the conjecture for any family of graphs which excludes a fixed minor (this includes planar graphs, for instance).

In Section 5, we modify a probabilistic decomposition of [17, 3] for use with growth-restricted graphs. Our modifications are two-fold. First, the parameters of our decomposition depend on ρ (and not on $n = |V|$ as in [17, 3]). This is essential to our application. Secondly, our decomposition is local in the sense that events which are far apart (in G) are mutually independent. As a result, we are able to apply the Lovász Local Lemma, yielding decompositions which, when combined with the results of Section 4, give $\dim(G) = O(\rho_G^3)$ for any graph G .

To obtain a tight upper bound of $\dim(G) = O(\rho_G \log \rho_G)$, we observe in Section 5.3 that the many steps of our embedding can be essentially performed “at once,” removing the need to amplify the individual probabilities at each step. Here, it is essential that every step of the embedding process is “local” with respect to its “scale.”

Finally, in Section 6, it is shown that all our results for $\dim(G)$ hold also for Linial’s variant $\dim_2(G)$ [15], yielding a conclusive answer to Question 2, and positively resolving the question of Benjamini and Schramm. This follows from a standard application of Chernoff-type tail bounds to the random processes employed in previous sections.

1.2 Related work

Notions of dimensionality for graphs were perhaps first considered by Erdős, Harary, and Tutte [8]. The geometric representations of graphs have been extensively studied in other settings; see, for instance, the survey of Lovász and Vesztergombi [18]. As mentioned before, the related study of low-distortion metric embeddings has received increasing attention in recent years (see the surveys [13, 19]).

Conjecture 1 is actually a dual of the bandwidth problem for graphs. The bandwidth problem asks for the minimum stretch of any edge in an embedding of the graph into $\mathbb{Z} = \mathbb{Z}_{\infty}^1$. Conjecture 1 asks for the minimum dimension needed to achieve a stretch of one (no stretch). Interestingly, the density bound $D = \max\{\frac{|B(v,r)|}{2^r}\}$ (a one-dimensional analogue of the growth rate), which is a straightforward lower bound on the bandwidth, is conjectured to be within a $\log |V|$ factor of the bandwidth (this gap is met, for instance, by expanders). From [9], we know that the bandwidth and density differ by only a $\text{polylog}(|V|)$ factor. However, the techniques employed for the bandwidth problem do not seem to help in resolving the dual question.

Imposing a growth restriction like $|B(v,r)| \leq r^\rho$ on a graph has many analogs in the analysis of

metric spaces, and in Riemannian geometry. For metric spaces, there are the notions of a *doubling metric* and *Ahlfors Q -regularity* (see [12]). A metric space (X, d) is called doubling if there exists a constant $L > 0$ such that every ball in X can be covered by L balls of half the radius. Assouad [2] showed that, for any fixed $0 < \epsilon < 1$, the metric space (X, d^ϵ) (with distances raised to the power ϵ) embeds into \mathbb{R}^k with distortion D , where k and D depend only on L . This theorem is similar in spirit to a number of the results presented here. Although Assouad's methods are significantly different and do not apply to Levin's problem, the relationship is not entirely superficial; the techniques of Section 5 were used in [11] to obtain a new (quantitatively almost optimal) proof of Assouad's theorem.

1.3 Preliminaries

For a point $x = (x_1, \dots, x_d) \in \mathbb{R}^d$, we write $\|x\|_2 = (\sum_{i=1}^d |x_i|^2)^{\frac{1}{2}}$ and $\|x\|_\infty = \max_{i=1}^d |x_i|$. For a graph G , we write $V(G)$ and $E(G)$ for the vertex and edge set of G , respectively.

Definition 1.1. For a graph $G = (V, E)$, a map $\varphi : V \rightarrow \mathbb{Z}^d$ is a contraction (or contractive) if $(u, v) \in E$ implies $\|\varphi(u) - \varphi(v)\|_\infty \leq 1$. Furthermore, if $\{\varphi_i\}$ is a finite set of mappings, we define the direct sum, $\varphi = \bigoplus_i \varphi_i$ to be the mapping $\varphi(u) = (\varphi_1(u), \varphi_2(u), \dots)$ (i.e., where coordinates are concatenated).

Notice that if a map $\varphi : V \rightarrow \mathbb{Z}^d$ is both contractive and injective, then G occurs as a subgraph of \mathbb{Z}_∞^d , and in particular, $\dim(G) \leq d$. We can think of any such embedding φ as consisting of d separate one-dimensional maps $\varphi_1, \dots, \varphi_d$ such that $\varphi = \bigoplus_{i=1}^d \varphi_i$. The following simple lemma will serve as our guide.

Lemma 1.2. Let $G = (V, E)$ and $\varphi = \bigoplus_{i=1}^d \varphi_i$ where $\varphi_i : V \rightarrow \mathbb{Z}$, then the following are true.

1. φ is a contraction \iff every φ_i is a contraction.
2. φ is injective \iff for every pair $u, v \in V$, there exists some φ_i such that $\varphi_i(u) \neq \varphi_i(v)$.

Unless otherwise stated, $\|\cdot\| = \|\cdot\|_\infty$ and all logarithms are to the base 2.

2 Trees

In this section we show that every tree T with growth rate ρ occurs as a subgraph of \mathbb{Z}_∞^d with $d = O(\rho)$ by exhibiting a map $\varphi : T \rightarrow \mathbb{Z}^d$ that is both contractive and injective.

2.1 Embedding trees by random walks

In light of Lemma 1.2, it is natural to define a distribution over random contractions and then argue that some such map must be injective.

Let $T = (V, E)$ be a tree whose growth rate is at most ρ . Fix an arbitrary root r of T , and let h be the height of T (so that $V \subseteq B(r, h)$). Define d_T to be the shortest path metric on T and let $c > 0$ be a sufficiently large constant to be determined later. Let $T_1, T_2, \dots, T_{c\rho}$ be $c\rho$ weighted copies of T , where the weight of every edge in T_i is chosen independently and uniformly at random from the set $\{-1, +1\}$. For each $v \in V$, let v_i be the sum of the edge weights on the unique path from r to v in T_i . Finally, define the image of v in $\mathbb{Z}^{c\rho}$ by $\varphi(v) = (v_1, v_2, \dots, v_{c\rho})$.

Clearly φ is a contraction. Now consider any two vertices $u, v \in V$ for which $d_T(u, v) \geq \sqrt{h}$. The probability that the images of u and v agree in any single coordinate, i.e. that $u_i = v_i$, is the probability that a random walk with $+1/-1$ steps and length \sqrt{h} ends at its starting point, namely $O(h)^{-1/4}$. Hence the probability that $\varphi(u) = \varphi(v)$ is at most $O(h)^{-c\rho/4}$. Observe that since T is contained in a ball of radius h centered at r , it contains at most h^ρ vertices. Taking a union bound over at most $h^{2\rho}$ pairs $\{u, v\} \in V^2$,

$$\Pr[\exists u, v \in V, d_T(u, v) \geq \sqrt{h} \text{ and } \varphi(u) = \varphi(v)] \leq h^{2\rho} O(h)^{-c\rho/4}.$$

It follows that there exists universal constants $h_0 > 0, c > 9$ such that for every tree T of height $h \geq h_0$, there exists a map $\varphi : V \rightarrow \mathbb{Z}^{c\rho}$ for which $d_T(u, v) \geq \sqrt{h} \implies \varphi(u) \neq \varphi(v)$. In what follows, we carefully utilize this simple but powerful observation to show that $\dim(T) = O(\rho)$ for any tree T , thus proving Conjecture 1 for the special case of trees.

2.2 Relative embeddings and rooted subtrees

Consider a tree $T = (V, E)$ with $\rho = \rho_T$ and fix a root r_T of T . Define a *rooted subtree* of T to be a connected vertex-induced subgraph X with a distinguished root r_X . Let $W = \{-1, 0, +1\}$ be the set of edge weights. For a rooted subtree $X = (V_X, E_X)$, we define a *d-dimensional relative embedding* of X to be a map $\mu_X : E_X \rightarrow W^d$. Finally, we will denote by $\mu_X^* : V_X \rightarrow \mathbb{Z}^d$ the (*absolute*) *embedding* induced by the relative embedding μ_X (with respect to the root r_X), which is the map obtained as follows: For a vertex $v \in V_X$, define its image $\mu_X^*(v) \in \mathbb{Z}^d$ as the sum of the edge weights $\mu_X(e)$ along the unique path from r_X to v in X . By our choice of W , induced embeddings are always contractions. Furthermore, given any contractive embedding $\varphi : V_T \rightarrow \mathbb{Z}^d$, there exists a unique *d-dimensional relative embedding* μ such that $\varphi = \mu^*$ (with respect to r_X). Let us define $\mathbf{0} = (0, 0, \dots, 0)$ to be the all-zero vector. Notice that the construction of Section 2.1, when applied to a subtree X , yields a relative embedding of X with the following desirable property.

Lemma 2.1 (Relative embeddings). *There exist constants h_0 and c such that for every rooted subtree $X = (V_X, E_X)$ of T with height at most h where $h \geq h_0$, there is a relative embedding $\mu_X : E_X \rightarrow W^{c\rho}$ such that, for all $u, v \in V_X$, $\mu_X^*(u) \neq \mu_X^*(v)$ whenever $d_T(u, v) \geq \sqrt{h}$.*

In essence, μ_X^* “separates” points in X which are far apart relative to the height of T . Notice that the above lemma only works for h sufficiently large. When h is bounded, i.e., $h = O(1)$, a brute force embedding suffices.

Lemma 2.2 (Small subtrees). *For any rooted subtree $X = (V_X, E_X)$ of T with height at most h_1 , there exists a relative embedding $\mu : E_X \rightarrow W^{\rho \log(h_1+2)}$ such that μ^* is injective.*

Proof. Let $V_X = \{v_1, v_2, \dots, v_m\}$ and define $\varphi(v_i) = B(i)$ where $B(i)$ is the binary representation of $i - 1$ written as a $\lceil \log m \rceil$ -dimensional vector. Now φ is clearly injective and also a contraction since $\|B(i) - B(j)\|_\infty \leq 1$ for all i, j . Finally, notice that since X is of height $h = O(1)$, we have $m \leq h^\rho$ and hence $\lceil \log m \rceil \leq \rho \log(h_1 + 2)$. Let μ be the unique relative embedding such that $\varphi = \mu^*$. It follows that $\mu : E_X \rightarrow W^{\rho \log(h_1+2)}$ is a relative embedding with μ^* injective. \square

Suppose $\{X_1, X_2, \dots, X_k\}$ is a collection of vertex-disjoint rooted subtrees of T , and let each $X_i = (V_i, E_i)$ have root r_i . Furthermore, suppose that for each X_i , we have a relative embedding $\mu_i : E_i \rightarrow W^d$. Then we can define a relative embedding μ on all of V by

$$\mu(e) = \begin{cases} \mu_i(e) & \text{if } e \in E_i \text{ for some } i \\ \mathbf{0} & \text{otherwise.} \end{cases}$$

Note that μ has the desirable property $\|\mu^*(u) - \mu^*(v)\| = \|\mu_i^*(u) - \mu_i^*(v)\|$ whenever $u, v \in V_i$. We will say that μ is obtained by *glueing* the relative embeddings $\{\mu_i\}$ together.

In the sequel, we will construct, for a given $\hat{h} > 0$ (which need not be the height of the tree T), an embedding $\varphi : V \rightarrow \mathbb{Z}^{O(\rho)}$ that satisfies $\varphi(u) \neq \varphi(v)$ whenever $\hat{h}^{1/2} \leq d(u, v) \leq \hat{h}$. (This is essentially the “single scale” version of the conjecture.) To do this, we will partition T into subtrees of height $O(\hat{h})$, find for each subtree a relative embedding that satisfies the desired property, and then glue all these relative embeddings into an embedding for T . There is the slight problem that for pairs u, v with $\hat{h}^{1/2} \leq d(u, v) \leq \hat{h}$ that end up in different subtrees, we have no guarantee that their images (under μ^*) will be distinct. To handle this, we will actually use two sets of disjoint subtrees which are “staggered” so that every pair u, v with $d(u, v) \leq h$ ends up in the same subtree in at least one of the sets. A far more challenging problem is that this embedding is guaranteed to handle only one value of h .

2.3 The leveled decomposition

Let $\text{diam}(T)$ be the diameter of T . Set $k = \lceil \log \log \text{diam}(T) \rceil$ and $\Delta = 2^{2^k}$, hence $\text{diam}(T) \leq \Delta \leq \text{diam}(T)^2$. We define k levels L_0, L_1, \dots, L_{k-1} as follows. Level i consists of two partitions of T into rooted subtrees. Denote these two partitions A_i and B_i and let $L_i = A_i \cup B_i$. The subtrees in L_i will cover T (in a sense that will be defined soon) and will each have height at most $3h(i)$, where $h(i) = \Delta^{1/2^i}$. (For convenience, we define $h(k) = 1$.) To form A_i , let O_i^A be the set of edges in T whose *depth* (i.e., distance from the root r_T) is a multiple of $3h(i)$. Removing O_i^A from T results in a collection of disjoint subtrees; let A_i consist of these subtrees, each rooted at its (unique) closest vertex to r_T . B_i is formed similarly, except that O_i^B is defined as the set of edges in T whose depth modulo $3h(i)$ is equal to $h(i)$ (rather than 0). The edges in O_i^A and O_i^B are called the *open edges* of level L_i . The next lemma is easy to verify. In particular, property (3) follows from the “staggering” of the two sets of subtrees A_i and B_i . Property (4) follows from the specifics of the decomposition; it provides a nesting that will turn out to be useful in Section 2.5.

Lemma 2.3 (The leveled decomposition). *For every tree $T = (V, E)$, the above construction satisfies the following properties.*

1. Each A_i and each B_i is a partition of V .
2. The height of any subtree $X \in L_i$ is at most $3h(i)$.
3. For any pair $u, v \in V$ with $d(u, v) \leq h(i)$, there is some $X \in L_i$ containing both u and v .
4. For every i , $O_{i+1}^A \subseteq O_i^A$; hence, each subtree in A_i is entirely contained in some subtree in A_{i+1} . The same holds for the subtrees in B_i . In this sense, each level is a refinement of the previous level.

Definition 2.4 (A separating map). *We say that $\varphi : V \rightarrow \mathbb{Z}^d$ separates A_i (and similarly B_i) if, for every $X \in A_i$ and for every pair $u, v \in V(X)$ with $h(i+1) \leq d(u, v) \leq h(i)$, we have $\varphi(u) \neq \varphi(v)$.*

Notice that if φ separates A_i and B_i for all $i \in \{0, 1, \dots, k-1\}$, then φ is injective (by the properties in Lemma 2.3).

2.4 A first attempt

Consider the partition A_i of T . Applying the embedding of Lemmas 2.1 and 2.2 to each $X \in A_i$ and glueing the relative embeddings together yields an induced embedding φ_i^A which separates A_i .

Let φ_i^B be a similar embedding obtained from the partition B_i , and let $\varphi_i : V \rightarrow \mathbb{Z}^{O(\rho)}$ be defined as $\varphi_i = \varphi_i^A \oplus \varphi_i^B$. Then φ_i separates A_i and B_i . Finally, the map $\varphi = \varphi_0 \oplus \dots \oplus \varphi_{k-1}$ separates every A_i and every B_i , and is hence injective (by Lemma 1.2). Since φ is also a contraction, it yields $\dim(T) = O(\rho k) = O(\rho \log \log \text{diam}(T))$. Unfortunately, this bound depends on the diameter of T and is therefore insufficient for our purposes.

2.5 Conserving randomness or “Not using all your ammo at once.”

In the preceding section, we used too many dimensions because we needed a distinct set of coordinates for every level. In essence, determining the embedding for level L_i leaves no randomness for “higher” levels $L_{i-1}, L_{i-2}, \dots, L_0$ (since all the edge weights in the relative embedding for L_i are determined).

Now consider the open edges of level L_i , namely, O_i^A and O_i^B , which run between disjoint subtrees. In Section 2.2, when the relative embeddings for subtrees are glued together, the open edges are assigned a weight of 0. But they might as well have been assigned any other weight in W^d . Clearly the resulting embedding would still be a contraction. Thus even after fixing a relative embedding for L_i , there is still some freedom left to us in deciding how to choose weights for the edges in O_i^A and O_i^B . It turns out that the randomness stored in the unassigned open edges is sufficient.

We will now show that, after finding a relative embedding for the subtrees in L_{i+2} , there is still enough randomness left to embed the subtrees in L_i simply by assigning random weights to the open edges of L_{i+2} . Notice that this process goes up two levels at a time, from L_{i+2} to L_i , so we will need to do it twice, once for “even” levels and once for “odd” levels. This will increase the number of coordinates used by only a factor of 2.

Theorem 2.5 (Embedding trees). *Any tree T with growth rate ρ occurs as a (not necessarily induced) subgraph of $\mathbb{Z}_\infty^{O(\rho)}$, thus $\dim(T) = O(\rho)$.*

Proof. We will construct four contractions, $\varphi_{\text{even}}^A, \varphi_{\text{odd}}^A, \varphi_{\text{even}}^B, \varphi_{\text{odd}}^B : V \rightarrow \mathbb{Z}^{O(\rho)}$. Let L_0, L_1, \dots, L_{k-1} be the levels of the decomposition for T , and assume for simplicity that k is odd. Then φ_{even}^A will separate $A_{k-1}, A_{k-3}, \dots, A_0$, φ_{odd}^A will separate $A_{k-2}, A_{k-4}, \dots, A_1$, and φ_{even}^B and φ_{odd}^B will satisfy similar properties for the B_i . It will follow from the discussion in Section 2.4, that $\varphi = \varphi_{\text{even}}^A \oplus \varphi_{\text{odd}}^A \oplus \varphi_{\text{even}}^B \oplus \varphi_{\text{odd}}^B$ is a contractive injection, providing the desired embedding of T into $\mathbb{Z}_\infty^{O(\rho)}$.

We will construct the map φ_{even}^A inductively. The other maps are constructed similarly. Let h_0 and c be the constant from Lemma 2.1 and let k_0 be the largest even integer such that $h(k_0) \geq h_0$. Since the trees in A_{k_0} have height at most $3h(k_0)$, and since we may assume without loss of generality that $c \geq \log(3h_0 + 2)$, Lemma 2.2 yields a relative embedding $\mu_{k_0} : E \rightarrow W^{c\rho}$ for which $\mu_{k_0}^*$ separates $A_{k-1}, A_{k-3}, \dots, A_{k_0}$.

The inductive step. Now assume that we have a relative embedding $\mu_{i+2} : E \rightarrow W^{c\rho}$ for which μ_{i+2}^* separates $A_{k-1}, A_{k-3}, \dots, A_{i+2}$. We will show the existence of a relative embedding $\mu_i : E \rightarrow W^{c\rho}$ which satisfies

1. For all $T \in A_{i+2}$ and all $u, v \in V(T)$, $\|\mu_{i+2}^*(u) - \mu_{i+2}^*(v)\| = \|\mu_i^*(u) - \mu_i^*(v)\|$;
2. μ_i^* separates A_i .

Since the subtrees of A_j for $j \geq i + 2$ are all completely nested within the subtrees of A_{i+2} (recall property (4) of Lemma 2.3), the first condition guarantees that μ_i^* separates $A_{k-1}, A_{k-3}, \dots, A_{i+2}$, since μ_{i+2}^* does.

To obtain μ_i from μ_{i+2} , we will only change the edge weights in O_{i+2}^A , i.e. those running between disjoint subtrees of A_{i+2} . Condition (1) then follows immediately. The construction of the relative embedding μ_i is actually probabilistic; we shall randomly change edge weights in O_{i+2}^A , and show that (2) is satisfied with positive probability. For every $e \in O_{i+2}^A$, choose $\mu_i(e)$ uniformly at random from $\{-1, +1\}^{c\rho}$. For all other edges e , define $\mu_i(e) = \mu_{i+2}(e)$.

Let us now show that with positive probability, μ_i^* separates A_i . Fix some $X \in A_i$ and consider two points $u, v \in V(X)$ such that $d(u, v) \geq h(i+1) = h(i)^{1/2}$. Let P_{uv} be the unique path from u to v in X . Since P_{uv} has length at least $h(i)^{1/2}$ and each subtree of A_{i+2} has height at most $3h(i+2) = 3h(i)^{1/4}$, P_{uv} must pass through at least $\frac{1}{3}h(i)^{1/4}$ such subtrees. In particular, the path includes at least $\frac{1}{3}h(i)^{1/4} - 2$ edges from O_{i+2}^A .

Now consider the part of P_{uv} which is composed of edges whose weights are already fixed (i.e., edges not in O_{i+2}^A). The sum of their weights is fixed, and the probability that a random walk of length at least $\frac{1}{3}h(i)^{1/4} - 2$ (along open edges) is equal to the negation of any fixed amount is at most $O(h(i))^{-1/8}$. This also upper bounds the probability that the images of u and v (under μ_i^*) agree in any single coordinate. So the probability of this occurring in $c\rho$ independent coordinates is

$$\Pr[\mu_i^*(u) = \mu_i^*(v)] = O(h(i))^{-c\rho/8}.$$

Finally, notice that X has height at most $3h(i)$, and thus contains at most $(3h(i))^{2\rho}$ pairs of vertices. Since $h(i) \geq h_0$ and we may assume that h_0 and c are sufficiently large constants (which are independent of i), the union bound $O(h(i))^{-c\rho/8}O(h(i))^{2\rho} < 1/2$ shows the existence of a map μ_i with the desired property in X . Continuing in this way for each disjoint subtree $X \in A_i$, we see that with positive probability, μ_i satisfies condition (2).

By induction, μ_0^* separates each of A_{k-1}, \dots, A_0 . Setting $\varphi_{\text{even}}^A = \mu_0^*$ completes the proof. \square

2.6 Alteration

We offer two simple lemmas on fixing maps which are sufficiently close to good embeddings.

Lemma 2.6 (Almost injective embeddings). *For a graph $G = (V, E)$, if there is a contractive map $\varphi : V \rightarrow \mathbb{Z}^d$ which is k -to-1, then there is a mapping $\varphi' : V \rightarrow \mathbb{Z}^{d+\lceil \log k \rceil}$ which is contractive and injective.*

Proof. To see this, suppose $z \in \text{Im}(\varphi)$ and let $\varphi^{-1}(z)_1, \dots, \varphi^{-1}(z)_k$ be the k possible preimages of z . Now define $\varphi'(\varphi^{-1}(z)_i) = (\varphi(z), B(i))$ where $B(i)$ is the $\lceil \log k \rceil$ -digit binary representation of $i - 1$. \square

Lemma 2.7 (Almost contractive embeddings). *For a graph $G = (V, E)$, if there is an injective mapping $\varphi : V \rightarrow \mathbb{Z}^d$ which satisfies $(u, v) \in E \implies \|\varphi(u) - \varphi(v)\|_\infty \leq k$, then there is a mapping $\varphi' : V \rightarrow \mathbb{Z}^{d(2+\log k)}$ which is contractive and injective.*

Proof. To get φ' , split \mathbb{Z}^d up into cubes of side length k . Now contract every such cube to its center, and then scale all the coordinates by a factor of $1/k$. It is easy to see that the resulting map is a contraction, but since each cube contained at most k^d points, the resulting map is only k^d -to-1. Applying Lemma 2.6, we can get a mapping which is injective and has $d + \lceil \log(k^d) \rceil$ dimensions. \square

2.7 Graphs without long induced simple cycles

For a graph G , let $\lambda(G)$ be the length of the longest induced simple cycle in G . We will use Lemmas 2.6 and 2.7 and the following theorem of Brandstädt, Chepoi, and Dragan [5, 6] to prove a result on graphs which have no long induced simple cycles.

Theorem 2.8 ([6]). *For any graph $G = (V, E)$, there exists a tree $T = (V, F)$ such that*

$$|d_G(u, v) - d_T(u, v)| = O(\lambda(G)). \quad (1)$$

Theorem 2.9. *For any graph G , let $\rho = \rho_G$, $\lambda = \lambda(G)$, then G occurs as a subgraph of $\mathbb{Z}_\infty^{O(\rho \log^2[\lambda+2])}$.*

Proof. Let $\rho = \rho_G$, $\lambda = \lambda(G)$, and let T be the corresponding tree of Theorem 2.8. First, setting $d_G(u, v) = 1$ in (1), we see that edges are stretched in T by at most $O(\lambda)$. Secondly, setting $d_T(u, v) = 1$ in (1), we see that edges in T correspond to paths in G of length at most $O(\lambda)$. It follows that

$$\rho_T = \sup_{x,r} \frac{\log |B_T(x, r)|}{\log r} \leq \sup_{x,r} \frac{\log |B_G(x, \lambda r)|}{\log r} \leq \sup_{x,r} \frac{\log(\lambda r)^\rho}{\log r} \leq \rho(1 + \log \lambda).$$

So we can embed T into $O(\rho \log(\lambda + 2))$ dimensions by Theorem 2.5. The same mapping is also an injective embedding for G that expands edges by at most $O(\lambda)$. Applying Lemma 2.7, we arrive at a contractive, injective embedding of G into $O(\rho \log^2[\lambda + 2])$ dimensions. \square

Corollary 2.10. *Conjecture 1 is true for any class of graphs in which $\lambda(G)$ is bounded, yielding $\dim(G) = O(\rho)$. This class includes trees and chordal graphs.*

3 Expanders

Before we jump into the proof of our main theorem, let us take a moment to prove a lower bound for expander graphs.

Definition 3.1 (Expander graphs). *An n -vertex graph $G = (V, E)$ will be called a $\Theta(k)$ -degree expander for $k = k(n) \geq 2$ if it has the following properties:*

- (a) *The degree of every vertex is $\Theta(k)$.*
- (b) *The diameter of G is $O(\log_k n)$.*
- (c) *Every two disjoint subsets of $n/\log n$ vertices are connected by a path of length $O(\log_k \log n)$.*

Observe that properties (b) and (c) follow from vertex expansion. Indeed, let $\Gamma(S)$ stand for the set of vertices with at least one neighbor in S , and suppose that for every $S \subset V$, we have $|\Gamma(S)| \geq \min\{\Omega(k|S|), \frac{2}{3}|V|\}$. Hence, there exists $t \leq O(\log_k n)$, such that every $u \in V$ satisfies $|\Gamma^t(u)| \geq \frac{2}{3}|V|$, and it follows that the diameter of G is at most $2t = O(\log_k n)$. Property (c) follows similarly.

Consequently, for every $3 \leq k(n) \leq \log n$ and every sufficiently large n , there exists a graph G satisfying properties (a)-(c). An even simpler way to obtain such a graph G is to take a (standard) 3-regular n -vertex expander H , and create G on the same vertex set by a connecting two vertices u, v whenever $d_H(u, v) \leq \log_3 k$. It is easy to satisfy property (a) by iteratively connecting by an edge the two vertices of lowest degree. A simple argument as above shows that the diameter of G is $O(\log n)$ and that every two sets of size $n/\log n$ are connected by a path of length $O(\log \log n)$. Properties (b) and (c) now follow from the fact that for very two vertices u, v we have $d_G(u, v) \leq O(d_H(u, v)/\log k)$.

3.1 A dimension lower bound

Theorem 3.2. *Let $G = (V, E)$ be a $\Theta(\log |V|)$ -degree expander, then $\dim(G) = \Omega(\rho_G \log \rho_G)$. In particular, Conjecture 1 is not true (for general graphs).*

Proof. Let $G = (V, E)$ be a $\Theta(k)$ -degree expander graph on n vertices (see Definition 3.1), with $1 \leq k \leq \log n$. It follows from Properties (a) and (b) that $\rho_G = \Theta(\frac{\log n}{\log \log_k n})$. We shall show that if G occurs as a subgraph of \mathbb{Z}_∞^d then $d \geq \Omega(\frac{\log n}{1 + \log \log_k \log n})$. Note that for $k = \log n$, this implies that $\dim(G) = \Omega(\log n) = \Omega(\rho_G \log \rho_G)$ and this lower bound is tight, up to constant factors, since the trivial upper bound $d = O(\log n)$ holds for any n -vertex graph (by a bijection into $\{0, 1\}^{\lceil \log n \rceil}$).

Assume for contradiction that G occurs as a subgraph of \mathbb{Z}_∞^d with $d = o(\frac{\log n}{1 + \log \log_k \log n})$. Let φ be the corresponding embedding of G into \mathbb{Z}_∞^d , and let φ_i be the projection of φ on the coordinate $i = 1, \dots, d$. Let the set S_i consist of the $n/\log n$ vertices $v \in V$ with smallest $\varphi_i(v)$, and let the set L_i consist of the $n/\log n$ vertices $v \in V$ with largest $\varphi_i(v)$, breaking ties arbitrarily.

We claim that $\varphi_i(V \setminus (S_i \cup L_i))$ is contained in an interval of size $O(\log_k \log n)$. Indeed, by property (c) above G contains a path of length $O(\log_k \log n)$ that connects some vertex $s \in S_i$ with some vertex $l \in L_i$, and since φ is contractive, $\varphi_i(l) - \varphi_i(s) \leq O(\log_k \log n)$. By the definition of S_i and L_i , for every $v \in V \setminus (S_i \cup L_i)$ we have $\varphi_i(s) \leq \varphi_i(v) \leq \varphi_i(l)$, which proves the claim.

Finally, the set of vertices $V' = V \setminus (\cup_{i=1}^d (S_i \cup L_i))$ contains at least $n - dn/\log n > n/2$ vertices. By the above claim, $\varphi(V')$ is contained in a subset of the lattice \mathbb{Z}^d formed by the Cartesian product of d intervals of size $O(\log_k \log n)$. However, this subset of \mathbb{Z}^d contains at most $(O(\log_k \log n))^d \leq n/2$ points, which contradicts the assumption that φ is injective. \square

3.2 A distortion lower bound

It is well-known that the bandwidth of an expander is $\Omega(n)$. The bandwidth may be seen as the maximum stretch of any edge in an injective embedding of the graph into \mathbb{Z}_∞^1 , thus embeddings into \mathbb{Z}_∞^d provide a generalization of the bandwidth. We show now that even in much higher dimensions, the edges of the expander must be stretched by a large factor.

Corollary 3.3. *For $1 \leq k \leq \log n$, any embedding of an n -vertex $\Theta(k)$ -degree expander G into $\mathbb{Z}_\infty^{O(\rho_G)}$ stretches at least one edge to length $(\log n)^{\Omega(1)}$.*

Proof. Proceed similar to the proof of Theorem 3.2, with $d = O(\rho_G) = O(\frac{\log n}{\log \log_k n})$. Now if every edge is stretched by at most α , then two sets of $n/\log n$ vertices have their lattice images within distance $O(\alpha \log_k \log n)$ of each other. It follows that $O(\alpha \log_k \log n)^{O(\rho_G)} \leq n/2$, and for $\alpha = (\log n)^{o(1)}$ we derive a contradiction. \square

4 Divide & conquer: Upper bounds from graph decomposition

In this section, we will use the ideas of Section 2 to prove a result on general graphs in terms of their decompositions (Theorem 4.2). As an example application of this general result, we will show that Conjecture 1 holds for graphs excluding a fixed minor (Section 4.4).

Outline. At a conceptual level, our embedding method for a general graph G bears some similarity to the case of trees, although several technical aspects differ significantly. We cannot, for example, form a coordinate by assigning edge weights independently at random (the weights

along every cycle would have to sum to 0). Here is an outline of the proof, with simplified notation and constants. Our first step is to focus on a single “scale” r , and show a contraction φ_r such that $\varphi_r(u) \neq \varphi_r(v)$ for every two vertices u, v with $r^{1/2} \leq d(u, v) \leq r$. The construction of φ_r follows a divide and conquer approach—we decompose the graph into (overlapping) “clusters”, embed each cluster separately, and then “glue” these embeddings together. The decomposition guarantees that for every pair u, v as above there exists at least one cluster C that contains them both, hence it suffices that the embedding φ_C of this cluster satisfies $\varphi_C(u) \neq \varphi_C(v)$. The decomposition further guarantees that each cluster C has diameter at most $r^{O(1)}$, and then the growth rate bound implies $|C| \leq r^{O(\rho_G)}$.

One key difference between trees and general graphs is in the “divide” stage. For trees, we were able to design such a decomposition directly (Section 2.3), using only two “layers”, i.e. two partitions of V . When embedding general graphs, we consider decompositions with more layers, but we must ensure that the number of layers is bounded by, say, $O(1)$ or $O(\rho_G)$, since it affects the dimension of the resulting embedding. Furthermore, for trees our decompositions were nested (in the sense that finer partitions were refinements of coarser ones); for general decompositions we will have to force this nesting property to hold.

Another key difference is in the “conquer” (or combining) stage. In trees, it is quite easy to glue embeddings of disjoint subtrees, using the concept of relative embeddings (Section 2.2). In general graphs, we ensure that embeddings of disjoint clusters can be glued together by restricting ourselves to cluster embeddings in which the “boundary” of the cluster is mapped to the all-zeros vector. One side effect of this restriction is that for $u, v \in C$ lying on or “close” to the boundary of C , we cannot require that $\varphi_C(u) \neq \varphi_C(v)$. In addition, to embed a single cluster C , we have to resort to a more sophisticated method, inspired by Rao [21]; roughly speaking, we employ another decomposition that breaks C into subclusters and map each subcluster independently at random. This inner decomposition of C into subclusters has the same requirements as the outer decomposition, but it is applied with a different parameter, namely—each subcluster’s diameter is less than $r^{1/2}$. To show that this embedding of C succeeds with high probability we use a union bound over the at most $|C|^2 \leq r^{O(\rho_G)}$ pairs $u, v \in C$. Notice that the inner decomposition guarantees, for pairs $u, v \in C$ as above, that $\varphi_C(u)$ and $\varphi_C(v)$ are independent, while the outer decomposition limits the size of the subproblem C , enabling the use of a union bound. On top of this, we have to adapt the technique of conserving randomness (Section 2.5) to this new embedding method.

Preliminaries. In what follows, let $G = (V, E)$ be a simple graph with growth rate $\rho = \rho_G$. A *cluster* of G is a simply subset $S \subseteq V$, though we will usually use this terminology in the context of a partition which contains S .

Define the *boundary of a cluster* S as $\partial S = \{u \in S : \exists (u, v) \in E, v \notin S\}$. The boundary of a collection \mathcal{C} of clusters is defined as $\partial \mathcal{C} = \cup_{S \in \mathcal{C}} \partial S$. For a cluster $S \subseteq V$ and a partition P of V , the *induced partition* (on the cluster S) is defined as $Q = \{C \cap S : C \in P\} \setminus \{\emptyset\}$. As before, let $\mathbf{0} = (0, 0, \dots, 0)$ be the all-zero vector.

For $u, v \in V$ let $d(u, v)$ be the distance between u and v in the shortest path metric of G . We stress that even when a particular cluster S is considered, $d(u, v)$ denotes the distance in G and not in S . In particular, define the *diameter* (sometimes called weak diameter) of $S \subseteq V$ to be $\text{diam}(S) = \sup_{u, v \in S} d(u, v)$. As usual, for $S \subseteq V$ we define $d(u, S) = \inf_{v \in S} d(u, v)$.

The padded decomposition. We first discuss our method of choosing clusters. For the rest of this section, fix an arbitrary constant $\alpha > 1$. We will not explicitly state the dependence of other constants on α , but it will be clear that $\alpha = O(1)$ suffices for our purposes.

Definition 4.1 (The padded decomposition). *A set $\{P_1, P_2, \dots, P_m\}$ of m partitions of V is called an r -padded decomposition of G with m layers if the following properties are satisfied.*

1. *If $C \in \cup_{i=1}^m P_i$, then $\text{diam}(C) \leq r^\alpha$.*
2. *For every $u \in V$ there exists some $C \in \cup_{i=1}^m P_i$ such that $B(u, 3r) \subseteq C$.*

We can now state our main result about embeddings obtained from graph decompositions. Its proof appears in Section 4.3.

Theorem 4.2 (Embedding via decomposition). *Let G be a graph with $\rho = \rho_G$. If for every $4 \leq r \leq \text{diam}(G)$ there exists an r -padded decomposition of G with m layers, then $\text{dim}(G) = O(m^2 \rho)$.*

4.1 Relative embeddings

Suppose we are given a cluster $S \subseteq V$. Define a d -dimensional relative embedding of S to be a contraction $\varphi : S \rightarrow \mathbb{Z}^d$ such that $\varphi(\partial S) = \mathbf{0}$, i.e. the boundary is mapped to $\mathbf{0}$. Suppose further that we would like to find a relative embedding of S with the following property (parameterized by $r > 0$): For every $u, v \in S$ with $d(u, v) > r$ and such that $B(u, 3r^{1/2}) \subseteq S$, we have $\varphi(u) \neq \varphi(v)$. In other words, since we are imposing the rather stringent condition that $\varphi(\partial S) = \mathbf{0}$, we only make requirements on vertices that are far enough from the boundary.

We will produce such an embedding using a technique inspired by the methods of Rao [21]. Each coordinate is formed by partitioning S into clusters of diameter at most r , so u, v as above must end up in different clusters. We then define the image of a vertex to be the distance from that vertex to the boundary of its cluster. To achieve injectiveness with high probability, we ‘‘perturb’’ the images by randomly contracting each cluster’s boundary inward.

For ease of notation, we define an r -inner decomposition to be an $r^{1/\alpha}$ -padded decomposition; in this case, clusters have $\text{diam}(C) \leq r$ and vertices have ‘‘padding’’ of the form $B(u, 3r^{1/\alpha})$. We now show how to use an r -inner decomposition to produce a good relative embedding.

Lemma 4.3 (Relative embeddings). *Suppose that G has an r -inner decomposition with m layers, and let $S \subseteq V$ be a cluster with $|S| \leq r^{O(\rho)}$. Then there exists a relative embedding $\varphi : S \rightarrow \mathbb{Z}^{O(m\rho)}$ such that for every $u, v \in S$ with $d(u, v) > r$ and $B(u, 3r^{1/\alpha}) \subseteq S$, we have $\varphi(u) \neq \varphi(v)$.*

Proof. The m partitions produced by the r -inner decomposition induce m partitions Q_1, \dots, Q_m of S (recall the definition of an induced partition). For each Q_j we will construct a map $\varphi_j : S \rightarrow \mathbb{Z}^{c\alpha\rho}$, where $c > 0$ is a constant to be determined later.

Fix some partition Q_j and form a single coordinate $\varphi_j^0 : S \rightarrow \mathbb{Z}$ as follows: For every $C \in Q_j$, choose some $r_C \in \{0, 1, \dots, r^{1/\alpha}\}$ uniformly at random and let $\partial_C^* = \{v \in C : d(v, \partial C) \leq r_C\}$ (this is the boundary of C randomly contracted inward). Now for each $u \in S$ let $C_u \in Q_j$ be the cluster containing u and define $\varphi_j^0(u) = d(u, \partial_{C_u}^*)$. Recall that $d(\cdot, \cdot)$ denotes distance in G , and thus $\varphi_j^0(u) = \max\{0, d(u, \partial_{C_u}^*) - r_{C_u}\}$.

Clearly $\varphi_j^0(u)$ is a contraction, since $(u, v) \in E$ implies that either u and v are in the same cluster $C \in Q_j$ and then $|d(u, \partial_C^*) - d(v, \partial_C^*)| \leq 1$, or each of them belongs to the boundary of its cluster and then $\varphi_j^0(u) = \varphi_j^0(v) = 0$. It is also clear that $u \in \partial S$ implies $u \in \partial C$ for some $C \in Q_j$ and hence $\varphi_j^0(u) = 0$. Thus $\varphi_j^0(\partial S) = 0$.

Now independently form $c\alpha\rho$ such coordinates (each time picking fresh values for the r_C) and let φ_j be the direct sum of the resulting maps, where $c > 0$ is a sufficiently large constant to be

determined later. Finally, set $\varphi = \varphi_1 \oplus \dots \oplus \varphi_m$. From the properties of φ_j , we conclude that $\varphi : S \rightarrow \mathbb{Z}^{cm\rho}$ is a contraction which maps ∂S to $\mathbf{0}$, i.e., a relative embedding.

Consider a pair u, v with $d(u, v) > r$ and such that $B(u, 3r^{1/\alpha}) \subseteq S$. It follows from property (2) of Definition 4.1 that there exists a partition Q_j of S and a subset $C \in Q_j$ for which $B(u, 3r^{1/\alpha}) \subseteq C$. Since $d(u, v) > r$, the two vertices u and v must lie in different subsets of Q_j . It follows that, in any single coordinate φ_j^0 of the map φ_j , the value of $\varphi_j^0(u)$ is distributed uniformly over an interval of size $r^{1/\alpha}$ independently of the value $\varphi_j^0(v)$, hence $\Pr[\varphi_j^0(u) = \varphi_j^0(v)] \leq r^{-1/\alpha}$. Thus the probability that u and v collide in all $c\alpha\rho$ coordinates of φ_j is $\Pr[\varphi_j(u) = \varphi_j(v)] \leq r^{-c\rho}$. Since $|S| \leq r^{O(\rho)}$, there are at most $r^{O(2\rho)}$ such pairs u, v , and hence the probability that there exists a pair that collides is at most $r^{O(2\rho)} r^{-c\rho} < 1/2$, if the constant c is chosen to be sufficiently large. The existence of a map φ satisfying the lemma follows. \square

4.2 A first attempt

Here is a simple approach which will fail in the end, but will give some intuition as to how the padded decomposition will be used. We will use the padded decomposition (Definition 4.1) to decompose G into layers of disjoint clusters. Using Section 4.1 we can then find a relative embedding for each cluster; glueing all these embedding together, we shall arrive at a good embedding for G . Note that the padded decomposition is being first to decompose the graph G into clusters which will be separately embedded, and then *inside* each cluster to compute a good relative embedding for that cluster.

Let $k = \lceil \log \log \text{diam}(G) \rceil$, and set $r_i = 2^{2^i}$ for $i \in \{1, \dots, k\}$, and $r_0 = 0$. We apply an r -padded decomposition with $r = r_1, \dots, r_k$. For each value of r , this decomposition will break the graph into clusters of diameter at most r^α such that every two vertices within a distance r are contained in some such cluster S and are “far” from the boundary of S (as otherwise we cannot ensure that they are “separated” by the relative embedding for S).

An embedding for one level. Assume that $i > 1$. Let $\{P_1, P_2, \dots, P_m\}$ be the partitions produced by the r_i -padded decomposition. We will show how to construct a contraction $\varphi_i : V \rightarrow \mathbb{Z}^{O(m^2\rho)}$ that satisfies: For every pair $u, v \in V$ with $r_{i-1} < d(u, v) \leq r_i$, we have $\varphi_i(u) \neq \varphi_i(v)$.

Fix a partition P_j of V . For every cluster $S \in P_j$, compute a relative embedding $\psi_S : S \rightarrow \mathbb{Z}^{O(m\rho)}$ by applying Lemma 4.3 with the parameter r set to $r_{i-1} = r_i^{1/2} \geq 2$. Note that the lemma is applicable since $\text{diam}(S) \leq r_i^\alpha$ implies $|S| \leq r_i^{\alpha\rho} = r_{i-1}^{O(\rho)}$. Now for every $u \in V$, set $\varphi_{ij}(u) = \psi_S(u)$ where $S \in P_j$ is the cluster containing u . Notice that this map is well-defined since P_j is a partition of V . Also, notice that it is a contraction, for suppose $(u, v) \in E$. If u and v are in the same cluster S , then $\|\varphi_{ij}(u) - \varphi_{ij}(v)\| = \|\psi_S(u) - \psi_S(v)\| \leq 1$ since ψ_S is a relative embedding, and hence a contraction. If u and v are in different clusters, then $\varphi_{ij}(u) = \varphi_{ij}(v) = \mathbf{0}$ since both of u and v are on the boundary of their cluster. Finally, set $\varphi_i = \varphi_{i1} \oplus \dots \oplus \varphi_{im}$.

Now consider some $u, v \in V$ with $r_{i-1} < d(u, v) \leq r_i$. By property (2) of Definition 4.1, there exists some partition P_j and a cluster $S \in P_j$ such that $B(u, 3r_i) \subseteq S$. Thus $u, v \in S$, and certainly $B(u, 3r_{i-1}^{1/\alpha}) \subseteq S$, so by Lemma 4.3, $\psi_S(u) \neq \psi_S(v)$. It follows that $\varphi_{ij}(u) \neq \varphi_{ij}(v)$, and hence $\varphi_i(u) \neq \varphi_i(v)$.

The base case. For $r = r_1 = O(1)$, we will construct φ_1 in a special way so that for all $u, v \in V$ with $0 < d(u, v) \leq r_1$ we have $\varphi_1(u) \neq \varphi_1(v)$. We break G into clusters using an r_1 -padded decomposition as above, but then revert to a much simpler relative embedding technique: Given a cluster $S = \{v_1, v_2, \dots, v_s\}$ with $\text{diam}(S) \leq r_1^\alpha$, define the relative embedding $\psi_S(v_i) = B(i)$ if $v_i \notin$

∂S and $\psi_S(v_i) = \mathbf{0}$ otherwise, where $B(i)$ is the binary representation of i as an $O(\rho)$ -dimensional vector. Notice that the number of coordinates meets our needs, since $s = |S| \leq r_1^{\alpha\rho} \leq 2^{O(\rho)}$. This map is a contraction and satisfies $\psi_S(u) \neq \psi_S(v)$ whenever $u, v \notin \partial S$. Using this technique in the above argument (instead of Lemma 4.3) yields the desired map φ_1 . In fact, this map uses only $O(m\rho)$ coordinates, so we append 0's to every image and extend it to $O(m^2\rho)$ coordinates.

Putting it all together. If we let $\varphi = \varphi_1 \oplus \varphi_2 \oplus \cdots \oplus \varphi_k$, we see that φ is a contractive, injective embedding since for any distinct $u, v \in V$, the distance $d(u, v)$ falls into some range $r_{j-1} < d(u, v) \leq r_j$ and thus $\varphi_j(u) \neq \varphi_j(v)$.

Assuming that for every $r \geq 4$ we can construct r -padded decompositions with m layers, then each φ_{ij} (which we obtained by applying Lemma 4.3) uses $O(m\rho)$ coordinates, and thus each φ_i uses $O(m^2\rho)$ coordinates. It follows that the final embedding φ uses $O(m^2\rho \log \log \text{diam}(G))$ coordinates in all. It turns out that this bound is of the right form, except for the dependence on $\text{diam}(G)$, so our next goal will be to eliminate this term. In the case of trees, we achieved this goal by exploiting some “untapped randomness”, namely, after fixing a relative embedding for a level, we were still free to assign arbitrary weights to the open edges of that level. In the next section, we exploit a similar observation, namely that the boundary of a cluster need not be mapped to $\mathbf{0}$, because the edges running between clusters are still “open.”

4.3 Forced nesting, contracted clusters, and untapped randomness

We improve over the preceding failed attempt by reusing the coordinates when proceeding inductively from finer partitions to coarser ones. Informally, the main idea is to consider every cluster of the finer partition as a single entity whose embedding is “rigid”, achieved by contracting each such cluster into a single vertex. For this approach to work, we need the padded decompositions to be nested, achieved by a “forced nesting” technique. We introduce these two notions and then prove Theorem 4.2.

Contracted clusters. Suppose we have a relative embedding ψ_C for each cluster C in a partition P . Previously, we “glued” these embeddings by setting $\Psi(u) = \psi_C(u)$ where $C \in P$ is the cluster containing u . This yielded a contraction Ψ defined on all of V with the property that whenever u, v belong to the same cluster $C \in P$, we have $\|\Psi(u) - \Psi(v)\| = \|\psi_C(u) - \psi_C(v)\|$. The following gives a simple way of maintaining this property, while allowing some freedom in choosing Ψ .

Definition 4.4 (Contracted graph). *Let P be a partition of V . The contracted graph (with respect to P) is the graph $\hat{G} = (\hat{V}, \hat{E})$ obtained from G by contracting, in the graph-theoretic sense, each cluster $C \in P$ to a single vertex.*

Each cluster $C \in P$ corresponds to a vertex in \hat{G} , and vice versa. Hence, we may identify $\hat{V} = P$ and then $\hat{E} = \{(C_1, C_2) : \exists (u_1, u_2) \in E, u_1 \in C_1 \in P, u_2 \in C_2 \in P\}$. We let $d_{\hat{G}}(\cdot, \cdot)$ be the shortest path metric in the contracted graph \hat{G} , and $B_{\hat{G}}(\cdot, \cdot)$ be a closed ball in \hat{G} . We shall keep using $d(\cdot, \cdot)$ and $B(\cdot, \cdot)$ when referring to the corresponding notions in the given graph G . A cluster \hat{S} in \hat{G} and its boundary $\partial\hat{S}$ are defined similarly to those in G .

Lemma 4.5. *Let $\hat{G} = (\hat{V}, \hat{E})$ be the contracted graph with respect to a partition P of V . Suppose that for each cluster $C \in P$ we have a relative embedding $\psi_C : C \rightarrow \mathbb{Z}^d$, and suppose that we have a contraction $\hat{\psi} : \hat{V} \rightarrow \mathbb{Z}^d$. For every $u \in V$, define the map $\Psi : V \rightarrow \mathbb{Z}^d$ as follows: $\Psi(u) = \hat{\psi}(C_u) + \psi_{C_u}(u)$, where $C_u \in P$ is the cluster containing u . Then Ψ is a contraction and for all u, v in the same cluster $C \in P$, we have $\|\Psi(u) - \Psi(v)\| = \|\psi_C(u) - \psi_C(v)\|$.*

Proof. By definition, for all $u, v \in C$ and $C \in P$, we have $\|\psi(u) - \psi(v)\| = \|\psi_C(u) - \psi_C(v)\|$. In particular, ψ contracts every edge whose endpoints are in the same cluster C . For $(u, v) \in E$ where u, v are in different clusters of P , we have $\|\psi(u) - \psi(v)\| = \|\hat{\psi}(C_u) - \hat{\psi}(C_v)\| \leq 1$ because ψ_P is a contraction and $(u, v) \in E$ implies $(C_u, C_v) \in \hat{E}$. \square

Forced nesting. Given two partitions P and Q of V , we say that P is a *refinement* of Q if for all $C \in P$ and there exists $S \in Q$ such that $C \subseteq S$. It turns out that we can force such a nesting and remove only a negligible amount of padding.

Lemma 4.6. *Let $\mathcal{P} = \{P_1, \dots, P_m\}$ be an r -padded decomposition of G , and $\tilde{\mathcal{P}} = \{\tilde{P}_1, \dots, \tilde{P}_m\}$ be an \tilde{r} -padded decomposition of G , where $r \geq \tilde{r}^\alpha$. Then we can modify \mathcal{P} so that, for every $j \in \{1, \dots, m\}$, the partition P_j is a refinement of \tilde{P}_j , with only a constant factor loss in the padding guarantee of Definition 4.1, namely condition (2) is replaced with:*

(2') For every $u \in V$ there exists some $C \in \cup_{i=1}^m P_i$ such that $B(u, 2r) \subseteq C$.

Proof. Whenever a cluster $C \in P_j$ contains only a portion of a cluster $C' \in \tilde{P}_j$ (i.e. $\emptyset \neq C \cap C' \neq C'$), modify P_j by breaking C into the cluster $C \setminus C'$ and a singleton cluster $\{u\}$ for every $u \in C \cap C'$. This process is continued (in an arbitrary order) until there is no such cluster C . When this happens, for every $j \in \{1, \dots, m\}$, the partition \tilde{P}_j is a refinement of P_j , as desired.

Notice that condition (1) of Definition 4.1 remains satisfied since each modification only decreases the diameter of clusters in P_j . Condition (2) is still satisfied if we replace $B(u, 3r)$ by $B(u, 3r - \tilde{r}^\alpha) \supseteq B(u, 2r)$, which completes the proof. \square

We are now ready to prove Theorem 4.2. Here is a brief outline: We shall construct relative embeddings from the bottom up, i.e., proceeding inductively from finer partitions to coarser partitions. Once we find an embedding that is “good” for level j , we modify it so that it becomes good also for level $j - t$, for some constant t . This modification will involve constructing a random embedding of the contracted graph (essentially via Lemma 4.3) and combining it with the existing embedding (using Lemma 4.5). Repeating this $t = O(1)$ times, starting at levels $1, 2, \dots, t$, respectively, will yield an embedding that uses only $O(tm^2\rho) = O(m^2\rho)$ coordinates.

Proof of Theorem 4.2 (Embedding via decomposition). Suppose that for every $4 \leq r \leq \text{diam}(G)$, and every cluster $S \subseteq V$, there exists an r -padded decomposition of S with m layers. Let $\alpha > 1$ be the constant from Definition 4.1. As before, define $k = \lceil \log \log \text{diam}(G) \rceil$, $r_i = 2^{2^i}$ for $i \in \{1, \dots, k\}$, and $r_0 = 0$. For every $i \in \{1, \dots, k\}$ let $\mathcal{P}_i = \{P_1^i, \dots, P_m^i\}$ be the r_i -padded decomposition of G . Let $t = t(\alpha)$ be a positive integer to be chosen later (e.g., $t \geq \alpha^3$ suffices assuming $\alpha \geq 2$).

We first make sure that this series of decompositions $\mathcal{P}_1, \dots, \mathcal{P}_k$ is nested. Iteratively, for $i = t + 1, t + 2, \dots, k$, apply Lemma 4.6 to the padded decompositions \mathcal{P}_i and \mathcal{P}_{i-t} . Thereafter, for every $j \in \{1, \dots, m\}$, the partition P_j^{i-t} is a refinement of P_j^i with only a constant factor loss in the padding guarantee (which will be sufficient for what follows).

Definition 4.7 (A separating map). *We say that $\varphi : V \rightarrow \mathbb{Z}^d$ separates a partition P_j^i if, for all $u, v \in V$ such that $r_{i-1} < d(u, v) \leq r_i$ and such that there is a cluster $S \in P_j^i$ with $B(u, 2r_i) \subseteq S$, we have $\varphi(u) \neq \varphi(v)$.*

To prove the theorem it suffices to construct, for each $j \in \{1, \dots, m\}$ and each $i_0 \in \{1, \dots, t\}$, a contractive map $\varphi_{i_0, j} : V \rightarrow \mathbb{Z}^{O(m\rho)}$ that separates every partition P_j^i for which $i \equiv i_0 \pmod{t}$. Indeed, the direct sum $\varphi = \bigoplus_{i_0, j} \varphi_{i_0, j}$ is a contractive embedding of G into the $O(tm^2\rho)$ -dimensional

lattice, which separates every P_i^j and is thus injective. Since t is a constant, it would follow that $\dim(G) = O(m^2\rho)$.

Fixing $j \in \{1, \dots, m\}$ and $i_0 \in \{1, \dots, t\}$, it remains to construct only the map $\varphi_{i_0, j}$ satisfying the stated properties. We shall actually prove, by induction on i , a more general assertion: If $c = c(t, \alpha)$ is a sufficiently large constant, then for every $i \in \{1, \dots, k\}$ with $i \equiv i_0 \pmod{t}$, there exists a contractive map $\psi^i : V \rightarrow \mathbb{Z}^{cm\rho}$ such that

- i. ψ^i separates each of $P_j^{i_0}, P_j^{i_0+t}, \dots, P_j^i$, and
- ii. the restriction of ψ^i to any cluster $C \in P_j^i$ is a relative embedding of C .

The base case. For $i = i_0$ we have $r_{i_0} \leq r_t = O(1)$, and thus we can use the base case from Section 4.2 to generate a map ψ^{i_0} that separates $P_j^{i_0}$. This map need only use $O(\rho)$ coordinates, since we have only a single partition $P_j^{i_0}$ at hand rather than m partitions, but we can extend the map to use $cm\rho$ coordinates by appending 0's to every image.

The inductive step. Suppose we have a contractive map ψ^i which satisfies properties (i) and (ii), and let us construct ψ^{i+t} using Lemma 4.5. Define $\hat{G} = (\hat{V}, \hat{E})$ as the contracted graph of G with respect to the partition P_j^i . For every cluster $C \in P_j^i$, let $\psi_C : C \rightarrow \mathbb{Z}^{cm\rho}$ be the restriction of ψ^i to C ; by the induction hypothesis, ψ_C is a relative embedding of C . It remains to define an embedding $\hat{\psi} : \hat{V} \rightarrow \mathbb{Z}^{cm\rho}$, and then we can let $\psi^{i+t} : V \rightarrow \mathbb{Z}^{cm\rho}$ be the map yielded by Lemma 4.5.

Since $\hat{V} = P_j^i$ is a refinement of P_j^{i+t} , the latter naturally yields a partition \hat{P}_j^{i+t} of \hat{V} , as follows. With every cluster $S \in P_j^{i+t}$ we associate a cluster $\hat{S} = \{C \in \hat{V} : C \subseteq S\}$ in \hat{G} . It is then easy to verify that $\hat{P}_j^{i+t} = \{\hat{S} : S \in P_j^{i+t}\}$ is a partition of \hat{V} .

Below, we shall define the mapping $\hat{\psi}$ on a single cluster $\hat{S} \in \hat{P}_j^{i+t}$ in \hat{G} . By construction, $\hat{\psi}$ will be a relative embedding of that cluster \hat{S} (with respect to \hat{G}) i.e. $\hat{\psi}(\partial\hat{S}) = \mathbf{0}$, and thus the resulting embedding will necessarily be a contractive embedding of the entire \hat{V} . In fact, we shall define $\hat{\psi}$ on each \hat{S} randomly.

Fix a cluster $S \in P_j^{i+t}$ and the corresponding $\hat{S} \in \hat{P}_j^{i+t}$. Now construct an ‘‘inner decomposition’’ for \hat{S} , as follows: Take an m -layer $r_{i+t}^{1/\alpha}$ -inner decomposition of G (note that $r_{i+t}^{1/\alpha^2} \geq 4$), and force P_j^i to be a refinement of each layer of this inner decomposition using Lemma 4.6. The forced nesting procedure is applicable because $r_{i+t}^{1/\alpha^2} \geq r_i^\alpha$. This procedure modifies only the inner decomposition (and not P_j^i), changing its padding guarantee to $2r_{i+t}^{1/\alpha^2}$. Each layer of the inner decomposition is a partition of V , and hence induces a partition of S . Denote these m partitions of S by Q_1, \dots, Q_m . Since P_j^i is a refinement of the inner decomposition, each partition Q_l of S naturally yields a partition of \hat{S} which we shall denote by \hat{Q}_l . (This is similar to the way \hat{P}_j^{i+t} was defined.)

We generate the map $\hat{\psi} : \hat{S} \rightarrow \mathbb{Z}^{cm\rho}$ in a random fashion, using an argument similar to Lemma 4.3.¹ Fix a partition Q_l and form a single coordinate $f : \hat{S} \rightarrow \mathbb{Z}$ randomly as follows: For every $C \in \hat{Q}_l$ choose a value $r_C \in \{0, 1, \dots, r_i\}$ uniformly at random and let $f(w) = \max\{0, d_{\hat{G}}(w, \partial\hat{Q}_l)\}$. Now independently form $cm\rho$ such coordinates (each time picking fresh values for the r_C), and let $\hat{\psi}_l : \hat{S} \rightarrow \mathbb{Z}^{cm\rho}$ be the direct sum of the resulting maps. We apply the above to every partition Q_l and set $\hat{\psi} = \hat{\psi}_1 \oplus \dots \oplus \hat{\psi}_m$.

¹Applying this lemma directly to \hat{S} would generate $\hat{\psi}$ such that $\hat{\psi}(C) \neq \hat{\psi}(C')$ for certain $C, C' \in \hat{V}$, but it does not guarantee that as a result $\psi^{i+t}(u) \neq \psi^{i+t}(v)$ for suitable $u \in C, v \in C'$.

Notice that once $\hat{\psi}$ is defined on \hat{S} , the map ψ^{i+t} yielded by Lemma 4.5 is defined on S . Since the former map is constructed in a random fashion, the latter mapping is randomized as well. The next two lemmas analyze this randomized embedding of S .

Claim 4.8. Fix $S \in P_j^{i+t}$. Then $\psi^{i+t} : S \rightarrow \mathbb{Z}^{c\alpha\rho}$ is a relative embedding of S .

Proof. Let us first show that $\hat{\psi} : \hat{S} \rightarrow \mathbb{Z}^{c\alpha\rho}$ is a relative embedding of \hat{S} (with respect to \hat{G}). Indeed, it is easy to see that every coordinate f generated as above using some partition \hat{Q}_l is a contraction, and that for every vertex $w \in \partial\hat{S}$, we have $w \in \partial\hat{Q}_l$ and hence $f(w) = 0$. It follows that $\hat{\psi}$, which is a direct sum of such maps f , is a relative embedding of \hat{S} .

Now consider ψ^{i+t} ; we know it is a contraction from Lemma 4.5, so we only need to show that $\psi^{i+t}(\partial S) = \mathbf{0}$. Fix $u \in \partial S$ and let $C_u \in P_j^i$ be the cluster containing u . Then $u \in \partial C_u$ (recall that $C_u \subseteq S$) and thus $\psi_{C_u}(u) = \mathbf{0}$. It also follows that $C_u \in \partial\hat{S}$, and hence $\hat{\psi}(C_u) = \mathbf{0}$. By definition, $\psi^{i+t}(u) = \hat{\psi}(C_u) + \psi_{C_u}(u) = \mathbf{0}$, which proves the claim. \square

Claim 4.9. Fix $S \in P_j^{i+t}$. Then with probability at least $1/2$, for all $u, v \in S$ with $r_{i+t-1} < d(u, v) \leq r_{i+t}$ and with $B(u, 2r_{i+t}) \subseteq S$, we have $\psi^{i+t}(u) \neq \psi^{i+t}(v)$.

Proof. Fix $u, v \in S$ with $r_{i+t-1} < d(u, v) \leq r_{i+t}$ and $B(u, 2r_{i+t}) \subseteq S$. Let $C_u \in P_j^i$ be the cluster containing u , and let $C_v \in P_j^i$ be similarly for v . Clearly, $u, v \in S \in P_j^{i+t}$; recalling that P_j^i is a refinement of P_j^{i+t} we get that $C_u, C_v \in \hat{S}$.

The ball $B = B(u, 2r_{i+t}^{1/\alpha^2})$ is contained in some cluster of the inner decomposition, even after the forced nesting with P_j^i . By the above, $B \subseteq S$ and thus B is entirely contained also in some cluster of some partition Q_l of S . Since clusters in P_j^i have diameter at most r^α (in G), we have:

- (a) $r_{i+t-1}/r_i^\alpha < d_{\hat{G}}(C_u, C_v) \leq r_{i+t}$.
- (b) $B_{\hat{G}}(C_u, 2r_{i+t}^{1/\alpha^2}/r_i^\alpha)$ is entirely contained in some cluster of \hat{Q}_l .

Now consider $\hat{\psi}_l(C_u)$ and $\hat{\psi}_l(C_v)$. Clusters in \hat{Q}_l have diameter (in \hat{G}) at most $r_{i+t}^{1/\alpha} \leq r_{i+t-1}/r_i^\alpha$, so (a) implies that C_u and C_v reside in different clusters of \hat{Q}_l . This, in conjunction with (b) (and because $2r_{i+t}^{1/\alpha^2}/r_i^\alpha \geq r_i$), implies that each coordinate of $\hat{\psi}_l(C_u)$ is distributed uniformly over an interval of size r_i , independently of the corresponding coordinate in $\hat{\psi}(C_v)$. Recalling that $\hat{\psi} = \hat{\psi}_1 \oplus \dots \oplus \hat{\psi}_m$, $\psi^{i+t}(u) = \psi^i(u) + \hat{\psi}(C_u)$, and $\psi^{i+t}(v) = \psi^i(v) + \hat{\psi}(C_v)$, we get that each of at least $c\alpha\rho$ coordinates of $\psi^{i+t}(u)$ is distributed uniformly over an interval of size r_i , independently of the corresponding coordinate in $\psi^{i+t}(v)$. Therefore,

$$\Pr[\psi^{i+t}(u) = \psi^{i+t}(v)] \leq r_i^{-c\alpha\rho}.$$

The number of pairs $u, v \in S$ is at most $|S|^2 \leq (r_{i+t}^\alpha)^{2\rho} = r_i^{2^{t+1}\alpha\rho}$, the claim follows via a union bound if only we choose the constant $c \geq 2^{t+2}$. \square

As mentioned before, we use the preceding claim to construct for every $S \in P_j^{i+t}$ a map $\hat{\psi} : \hat{S} \rightarrow \mathbb{Z}^{O(m\rho)}$. This collection of maps gives an embedding $\hat{\psi} : \hat{V} \rightarrow \mathbb{Z}^{O(m\rho)}$, and using Lemma 4.5 we produce from it an embedding $\psi^{i+t} : V \rightarrow \mathbb{Z}^{O(m\rho)}$ that satisfies the induction hypothesis for $i+t$, and this completes the proof of Theorem 4.2. \square

4.4 Graphs excluding a fixed minor

Let G be a graph that excludes a $K_{s,s}$ minor for some fixed s . By adapting a decomposition technique of Klein, Plotkin, and Rao [14] we construct, for any value $r \geq 1$, an r -padded decomposition of G with only $O(2^s)$ layers. Applying Theorem 4.2, we then arrive at the main result of this section.

Theorem 4.10 (Excluded minor families). *Conjecture 1 is true for any family of graphs that excludes a fixed minor. For such graphs, $\dim(G) = O(\rho_G)$.*

Proof. By Theorem 4.2 it suffices to show how to produce an r -padded decomposition with $m = 2^s$ layers for any graph that excludes a $K_{s,s}$ minor, as this implies that $\dim(G) \leq O(4^s \rho_G)$.

To this end, consider such a graph $G = (V, E)$ and fix a value r . First, construct a Breadth-First-Search (BFS) tree from an arbitrary vertex $v \in V$, and compute for every vertex its BFS level (i.e., its distance from v). Then cut the tree every $12r$ BFS levels by removing any edge connecting a vertex of BFS-level j to a vertex of BFS-level $j + 1$ for any $j \equiv 0 \pmod{12r}$. Let C_0 denote the resulting set of connected components of G . Next, take another copy of G and cut it similarly but starting from BFS level $6r$, i.e., remove any edge connecting a vertex of BFS-level j to a vertex of BFS-level $j + 1$ for any $j \equiv 6r \pmod{12r}$. Let C_1 denote the resulting set of connected components of G . Now for each C_i , apply the same procedure on all the connected components in C_i , namely, choose an arbitrary vertex, construct a BFS tree, and form two sets of connected components C_{i0} and C_{i1} by making the cuts as above (staggered, each at intervals of size $12r$). Repeat this process s times and let C_q for $q \in \{0, 1\}^s$ denote the 2^s final sets of connected components. From [14] we know that every connected component in every final C_q has diameter at most $O(r)$. In addition, for every vertex u the entire ball $B(u, 3r)$ is uncut in at least one final set C_q , because each step performs two different sets of “staggered” cuts, at least one of which must avoid the entire ball $B(u, 3r)$. \square

5 A general dimension upper bound

In this section we give a tight upper bound on the dimension of general graphs: $\dim(G) = O(\rho_G \log \rho_G)$ for any graph G . (In Section 3, we showed that this upper bound is met by expanders.) First, we devise a decomposition for growth-restricted metrics (Section 5.1) and use Theorem 4.2 to obtain a weaker upper bound of $O(\rho_G^3)$ (Section 5.2). Then, by combining the previous arguments more carefully and utilizing some Chernoff-type tail bounds, we obtain the aforementioned tight upper bound (Section 5.3). We shall use some terminology from Section 5.2.

5.1 Partitioning of growth-restricted graphs

Linial and Saks [17] and Bartal [3] show that for any graph $G = (V, E)$ and $1 \leq r \leq \text{diam}(G)$, there exists a probabilistic partitioning of G into disjoint clusters of diameter at most $O(r \ln |V|)$, such that for any pair of vertices $u, v \in V$, the probability that u and v end up in different clusters is at most $d(u, v)/r$. Let $\rho = \rho_G$. In this section, we give a similar decomposition, but we replace the diameter bound of $O(r \ln |V|)$ with a bound that is independent of $|V|$, namely $O(\rho r \ln r)$, for any $r \geq \rho$. Our partitioning method is similar to those of [17] and [3], but different in a subtle and crucial way: It is local. Events which are sufficiently far apart are mutually independent.

First, take the continuous exponential distribution with mean $r > 0$, truncate it at some value $M > 0$ and rescale the remaining density function. The resulting distribution, which we denote $\text{Texp}(r, M)$, has density function $p(z) = \frac{e^{M/r}}{r(e^{M/r}-1)}e^{-z/r}$ for $z \in (0, M)$.

The partitioning procedure. Let $V = \{v_1, v_2, \dots, v_n\}$ and let $r \geq \rho$. For each $v_t \in V$, choose independently a radius r_t according to the distribution $\text{Texp}(r, 8\rho r \ln r)$. Now define $S_t = B(v_t, r_t) \setminus \cup_{i=1}^{t-1} B(v_i, r_i)$ as the set of vertices v for which $B(v_t, r_t)$ is the first ball containing v . Finally, define the set of clusters to be $\mathcal{C} = \{S_1, \dots, S_n\}$.

It is easy to see that \mathcal{C} is a partition of V , and that the (weak) diameter of every cluster $C \in \mathcal{C}$ is bounded by $\text{diam}(C) \leq 16\rho r \ln r$. Further analysis will require the following simple facts. In particular, (3) shows that if $M \geq 2r$, the truncated exponential distribution is ‘‘almost’’ memoryless.

Fact 5.1. Consider a random variable $R \sim \text{Texp}(r, M)$ for $M \geq 2r > 0$. Then,

1. For all $\beta \geq 0$, $\Pr[R \geq \beta] \leq 2e^{-\beta/r}$.
2. For all $\beta \geq 0$, $\Pr[R \leq \beta] \leq 2(1 - e^{-\beta/r}) \leq 2\beta/r$.
3. For all $\beta \geq 0$ and $R_0 \leq M/2$, $\Pr[R \leq R_0 + \beta \mid R \geq R_0] \leq 2\beta/r$.

For a vertex $u \in V$ and $x \geq 0$, let \mathcal{E}_u^x be the event that $B(u, x)$ is split between multiple clusters, i.e., that no cluster $C \in \mathcal{C}$ fully contains $B(u, x)$.

Lemma 5.2. Let $u \in V$ and $r \geq 16\rho$, and $x \geq 0$. Then $\Pr[\mathcal{E}_u^x] \leq 10x/r$.

Proof. Assume $x \leq r$ (the theorem says nothing for larger x) and let $\mathcal{B} = B(u, x)$, $B_t = B(v_t, r_t)$. Let us say that the ball \mathcal{B} is cut by the ball B_t if $\emptyset \neq S_t \cap \mathcal{B} \neq \mathcal{B}$ while for all $i < t$, $S_i \cap \mathcal{B} = \emptyset$. \mathcal{E}_u^x is precisely the event that there is a ball B_t that cuts \mathcal{B} . Let us separate these balls B_t into two classes, depending on the distance from v_t to u . Define \mathcal{E}_{far} to be the event that there exists B_t that cuts \mathcal{B} and $d(v_t, u) \geq 4\rho r \ln r$. Define $\mathcal{E}_{\text{near}}$ to be the event that there exists B_t that cuts \mathcal{B} and $d(v_t, u) < 4\rho r \ln r$.

Fix v_t with $d(v_t, u) \geq 4\rho r \ln r$ and notice that by Fact 5.1,

$$\Pr[B_t \text{ cuts } \mathcal{B}] \leq \Pr[r_t \geq 4\rho r \ln r - x] \leq 2r^{-4\rho} e^{x/r} \leq 6r^{-4\rho}.$$

But the number of such v_t for which B_t can possibly cut \mathcal{B} is at most the number of vertices in a ball of radius $8\rho r \ln r + x \leq r^3$ which is at most $r^{3\rho}$. Taking a union bound over all such possible v_t , we see that $\Pr[\mathcal{E}_{\text{far}}] \leq 6r^{-4\rho} r^{3\rho} \leq 6/r^\rho \leq 6/r$. Thus we are left only to bound the probability of $\mathcal{E}_{\text{near}}$.

Let the random variable T be the minimum t such that $B_T \cap \mathcal{B} \neq \emptyset$ (note that possibly $v_T \in \mathcal{B}$). The ball B_T can either cut \mathcal{B} (in which case \mathcal{E}_u^x occurs) or contain \mathcal{B} (and then $\mathcal{B} \subseteq S_T$ is not cut by any ball B_t). By the principle of deferred decision it suffices to upper bound the conditional probability $\Pr[\mathcal{E}_{\text{near}} \mid T = t]$ for an arbitrary t . To this end, we may assume that $d(v_t, u) \leq 4\rho r \ln r$ (as otherwise this conditional probability is 0) and then $\mathcal{E}_{\text{near}}$ happens if and only if B_t cuts \mathcal{B} , which in turn happens only if $r_t \leq d(v_t, u) + x$. Hence,

$$\Pr[\mathcal{E}_{\text{near}} \mid T = t] \leq \Pr[r_t \leq d(v_t, u) + x \mid r_t \geq d(v_t, u) - x] \leq \frac{4x}{r},$$

where we have used Fact 5.1 in conjunction with $d(v_t, u) \leq 4\rho r \ln r$. Thus, $\Pr[\mathcal{E}_{\text{near}}] = \sum_t \Pr[T = t] \cdot \Pr[\mathcal{E}_{\text{near}} \mid T = t] \leq 4x/r$ and $\Pr[\mathcal{E}_u^x] \leq \Pr[\mathcal{E}_{\text{near}}] + \Pr[\mathcal{E}_{\text{far}}] \leq 10x/r$. \square

5.2 Layered decomposition of growth-restricted graphs

Now we describe how to obtain an r -padded decomposition with $O(\rho_G)$ layers for general graphs G . Plugging these values into Theorem 4.2 yields an embedding into $O(\rho_G^3)$ dimensions. We will only be able to show the existence of such decompositions under the assumption that $r \geq \rho$. In the case where $r \leq 16\rho$, clusters of diameter $r^{O(1)}$ have at most $\rho^{O(\rho)}$ points, so we will be able to embed these by brute force using only $O(\rho \log \rho)$ dimensions (similar to the base case of Section 4.2). The final result appears in Theorem 5.5.

Theorem 5.3 (Decomposition theorem). *For every graph $G = (V, E)$ and every $r \geq 16\rho_G$, there exists an r -padded decomposition with $m = O(\rho_G)$ layers.*

Proof. Let $\rho = \rho_G$ and assume $r \geq 16\rho$. To produce a single layer of the decomposition (a partition of V into clusters), we will use the procedure of Section 5.1, with the parameter r (in that procedure and in Lemma 5.2) set to r^2 . Notice that the clusters produced have diameter at most $32\rho r^2 \ln r \leq r^4$. For a vertex $v \in V$, let \mathcal{E}_v be the event that the ball of radius $3r$ about v is cut (i.e., split amongst two or more clusters). From Lemma 5.2, we know that $\Pr[\mathcal{E}_v] \leq 3/r$.

Now produce m layers independently (with fresh random coins each time) and let \mathcal{E}_v^m be the event that the ball of radius $3r$ about v is cut in *every* layer. Clearly $\Pr[\mathcal{E}_v^m] \leq (3/r)^m$. We would like to say that $\Pr[\bigwedge_{v \in V} \overline{\mathcal{E}_v^m}] > 0$. If we could show this with $m = O(\rho)$, the theorem would follow. And indeed, this is our goal. We will employ the following symmetric form of the Lovász Local Lemma, see e.g. [1].

Lemma 5.4 (Lovász Local Lemma). *Let A_1, \dots, A_n be events in an arbitrary probability space. Suppose that for each A_i there is a set that contains all but at most d of the other events A_j , such that A_i is mutually independent of this set of events. If for all $i \in \{1, \dots, n\}$ we have $\Pr[A_i] \leq p$, and $ep(d+1) \leq 1$, then $\Pr[\bigwedge_{i=1}^n \overline{A_i}] > 0$.*

Let $r_1 = 2r^3 \ln r + 6r$. An event \mathcal{E}_u^m is mutually independent of all events \mathcal{E}_v^m for which $d(u, v) > r_1$ because every ball in the partitioning of Section 5.1 has radius at most $r^3 \ln r$ and thus cannot intersect both $B(u, 3r)$ and $B(v, 3r)$. It follows that \mathcal{E}_u^m is mutually independent of the set of all events \mathcal{E}_v^m except those for which $v \in B(u, r_1)$, and there are at most $d = r_1^\rho$ such vertices v . Thus if $\Pr[\mathcal{E}_u^m] \leq 1/e(d+1)$, we can apply the local lemma and the theorem is proved. But this is easily accomplished by choosing say $m = \lceil 8\rho \rceil$. By applying Lemma 5.4 we conclude that there exists an r -padded decomposition for V (with $\alpha = 4$). \square

Theorem 5.5. *For every graph G with growth rate ρ_G , $\dim(G) = O(\rho_G^3)$.*

Proof (sketch). We provide only a sketch of the proof, since a better upper bound is given in the next section. We use the proof of Theorem 4.2, except that instead of the induction's base case being $r = O(1)$, we start with a level corresponding to $r = \rho^{O(1)}$. In this case, clusters have diameter $r^{O(1)} = \rho^{O(1)}$, so we can easily give a relative embedding for each cluster using only $O(\rho \log \rho)$ coordinates, similar to the base case of 4.2. The rest of the proof then proceeds unchanged, including the inductive step in which we use the decomposition provided by Theorem 5.3. \square

Remark: Algorithmization of the Local Lemma. Although most of the techniques in this paper can easily be interpreted algorithmically, this is not immediately obvious for our use of the local lemma. Fortunately, it is not difficult to see that standard techniques suffice (see, e.g., [1, Chapter 5]).

5.3 A tight upper bound

As mentioned previously, Theorem 5.3, combined with Theorem 4.2, shows that $\dim(G) = O(\rho_G^3)$ for every graph G . By carefully combining the previous arguments and utilizing some Chernoff-type tail bounds, we are able to find a tight upper bound, $\dim(G) = O(\rho_G \log \rho_G)$; see Theorem 5.8 below.

We first strengthen the decomposition of Theorem 5.3. Given m layers (partitions of V) P_1, P_2, \dots, P_m , we say that a vertex $u \in V$ is *padded* in a layer j if there exists a cluster $C \in P_j$ such that $B(u, 3r) \subseteq C$ (otherwise, we say that u is *unpadded* in layer j). We next show a decomposition in which every vertex is padded in most of the layers (rather than in one layer).

Theorem 5.6 (Strengthened decomposition theorem). *For every graph $G = (V, E)$ and every $r \geq 36\rho_G$, there exists an r -padded decomposition with $m = O(\rho_G)$ layers, in which:*

(2'') *For every $u \in V$ there are $\frac{3}{4}m$ partitions P_j in which there is $C \in P_j$ with $B(u, 3r) \subseteq C$.*

Proof. Similar to the proof of Theorem 5.3, we construct $m = O(\rho_G)$ layers of randomized partitions that always satisfy requirement (1), and argue that, with positive probability, requirement (2'') is satisfied. The probability that u is unpadded in a single layer is at most $3/r$ (this followed from Lemma 5.2), so the expected number of layers in which u is unpadded is at most $3m/r$. We now need the following Chernoff-type tail bound (see, e.g., [20, Chapter 4]).

Lemma 5.7 (A tail bound). *Let X_1, X_2, \dots, X_n be independent Poisson trials such that, for $1 \leq i \leq n$, $\Pr[X_i = 1] = p_i$ and $0 < p_i < 1$. Then for $X = \sum_i X_i$, $\mu = E[X]$, and any $\delta > 0$,*

$$\Pr[X > (1 + \delta)\mu] < \left(\frac{e^\delta}{(1 + \delta)^{1+\delta}} \right)^\mu < \left(\frac{e}{1 + \delta} \right)^{(1+\delta)\mu}.$$

Let m_u be the expected number of layers in which a vertex u is unpadded, and let \mathcal{E}_u be the event that u is unpadded in more than $\frac{1}{4}m$ layers. Then, applying the above lemma,

$$\Pr[\mathcal{E}_u] = \Pr \left[Y_u > \frac{r}{12} \cdot \frac{3m}{r} \right] \leq \frac{1}{(r/12e)^{m/4}}.$$

Applying Lemma 5.4 the same way we did in the proof of Theorem 5.3 for a suitable $m = O(\rho_G)$, we conclude that $\Pr \left[\bigwedge_{v \in V} \overline{\mathcal{E}_v}^m \right] > 0$. \square

Theorem 5.8 (Embedding of general graphs). *For every graph $G = (V, E)$ with growth rate ρ_G , $\dim(G) = O(\rho_G \log \rho_G)$.*

Proof. We adapt the proof of Theorem 4.2, as follows. First, we use the strengthened decomposition from Theorem 5.6. Second, we employ a more careful analysis that exploits the independence of coordinates constructed from different layers together with the local lemma, instead of applying a union bound separately in each cluster.

In the sequel, we describe the modifications to that proof. In fact, we will only show that any one level (in the sense of Section 4.2) can be embedded into $O(\rho \log \rho)$ dimensions. Using the nesting techniques of Section 4.3, the existence of a contractive and injective embedding that uses only $O(\rho \log \rho)$ coordinates follows.

Let $\rho = \rho_G$. Fix $r = 2^{2^i}$ and suppose we are given an r -padded decomposition with $m = O(\rho)$ layers P_1, P_2, \dots, P_m strengthened as per Theorem 5.6. We may assume further that $m \geq \rho$,

because otherwise we can just duplicate every layer $\lceil \rho/m \rceil$ times. Let $c > 0$ be a constant to be determined later. We shall construct, for each cluster $S \in P_j$, a relative embedding $\psi_S : S \rightarrow \mathbb{Z}^{c \log \rho}$, such that for all $u, v \in V$ with $\sqrt{r} < d(u, v) \leq r$ there exist a layer P_j and a cluster $S \in P_j$ such that $u, v \in S$ and $\psi_S(u) \neq \psi_S(v)$. Letting $\varphi_j(u) = \psi_{S_u}(u)$ where $S_u \in P_j$ is the cluster containing u , and setting $\varphi = \bigoplus_{j=1}^m \varphi_j$, we will conclude that $\varphi(u) \neq \varphi(v)$ for all $u, v \in V$ with $\sqrt{r} < d(u, v) \leq r$. As before, in the base case we shall replace the requirement $\sqrt{r} < d(u, v) \leq r$ with $0 < d(u, v) \leq r$.

The base case. Assume $16\rho \leq r \leq \rho^{O(1)}$ (note that this is where our decomposition breaks down). We produce, for every cluster $S \in P_j$, a relative embedding $\psi_S : S \rightarrow \mathbb{Z}^{c \log \rho}$ as follows: If $u \notin \partial S$, then let $\psi_S(u)$ be a $(c \log \rho)$ -dimensional vector chosen uniformly at random from $\{0, 1\}^{c \log \rho}$, and let $\psi_S(u) = \mathbf{0}$ otherwise. This ψ_S is clearly a contraction.

Consider two vertices u, v with $0 < d(u, v) \leq r$ and let P_j be a layer in which u is padded. In this layer, u and v belong to the same cluster $S \in P_j$ and $u \notin \partial S$, so $\Pr[\psi_S(u) = \psi_S(v)] \leq (\frac{1}{2})^{c \log \rho} = \rho^{-c}$. Let $\mathcal{E}_{u,v}$ be the event that, in the resulting embedding $\varphi = \bigoplus_{j=1}^m \varphi_j$, we have $\varphi(u) = \varphi(v)$. For this event to happen, it must be that in every layer in which u is padded, we have $\psi_S(u) = \psi_S(v)$. It then follows that

$$\Pr[\mathcal{E}_{u,v}] = \Pr[\varphi(u) = \varphi(v)] \leq \rho^{-cm/2} \leq \rho^{-c\rho/2}.$$

There are $\Omega(|V|)$ events $\mathcal{E}_{u,v}$ and we would like to argue that with positive probability, none of them occur. Again, the local lemma comes to our rescue. It is not difficult to see that $\mathcal{E}_{u,v}$ is independent of all events $\mathcal{E}_{u',v'}$ for which $d(u, u') > r$ (because the image of u is chosen independently of the images of all the corresponding v, u' , and v'). It follows that $\mathcal{E}_{u,v}$ is mutually independent of all but at most $d = r^{2\rho} \leq \rho^{O(\rho)}$ other events. Choosing the constant c to be a large enough relative to the constants in the bound $r \leq \rho^{O(1)}$, we see that $\Pr[\mathcal{E}_{u,v}] \leq \rho^{-c\rho/2} \leq 1/e(d+1)$. Thus, applying Lemma 5.4 yields an embedding for which none of the events $\mathcal{E}_{u,v}$ occur.

Higher levels. Assume $r \geq (16\rho)^2$. Recall that in Theorem 4.2, for each layer P_j and each cluster $S \in P_j$, we produced a random relative embedding $\psi_S : S \rightarrow \mathbb{Z}^{O(m \log \rho)}$ by constructing (using Lemma 4.3) $O(\log \rho)$ coordinates from each layer of an m -layer $r^{1/2}$ -inner decomposition. Here, instead, for each $S \in P_j$ we shall produce a random relative embedding $\psi_S : S \rightarrow \mathbb{Z}^c$ (which is of course stronger than using $O(\log \rho)$ coordinates), by constructing only $c = O(1)$ coordinates from layer j of the inner decomposition. More formally, take an m -layer $r^{1/2}$ -inner decomposition of G . Now for each P_j and each cluster $S \in P_j$, construct each of the c coordinates of ψ_S as follows (this is analogous to Lemma 4.3): Let Q_j be the partition of S induced by layer j of the inner decomposition. For every cluster $C \in Q_j$, choose $r_C \in \{0, 1, \dots, r^{1/2\alpha}\}$ uniformly at random. For every $u \in S$, let $C_u \in Q_j$ be the cluster containing u , and define the image of u to be $\max\{0, d(u, \partial C_u) - r_{C_u}\}$. Clearly, ψ_S is a relative embedding of S .

Consider two vertices u, v with $\sqrt{r} < d(u, v) \leq r$. Since u is padded in at least $\frac{3}{4}m$ layers each decomposition (i.e., the layers $\{P_1, \dots, P_m\}$ and in the inner decomposition), we see that for at least $\frac{1}{2}m$ values of j , the vertex u is padded both in P_j and in layer j of the inner decomposition. Fix such j , and let $S \in P_j$ be the cluster containing u . It follows that u, v belong to the same cluster $S \in P_j$, but to different clusters of the inner decomposition. Since u is padded in layer j of the inner decomposition, each coordinate of $\psi_S(u)$ is chosen at random from an interval of size at least $r^{-1/2\alpha}$, independently of $\psi_S(v)$, and thus the probability it collides with the corresponding coordinate of $\psi_S(v)$ is at most $r^{-1/2\alpha}$. Hence,

$$\Pr[\varphi_j(u) = \varphi_j(v)] = \Pr[\psi_S(u) = \psi_S(v)] \leq r^{-c/2\alpha},$$

and the embedding $\varphi = \bigoplus_{j=1}^m \varphi_j$ satisfies $\Pr[\varphi(u) = \varphi(v)] \leq r^{-cm/4\alpha} \leq r^{-c\rho/4\alpha}$. Finally, we would like to apply Lemma 5.4 on the events $\mathcal{E}_{u,v} = \{\varphi(u) = \varphi(v)\}$ where $\sqrt{r} < d(u,v) \leq r$. It can be seen that every event $\mathcal{E}_{u,v}$ is mutually independent of all the other events $\mathcal{E}_{u',v'}$ but the $r^{2\rho}$ events for which $d(u,u') \leq 3r$ (because every cluster of the inner decomposition is mapped independently). Hence, for $c > 0$ a sufficiently large constant we can apply Lemma 5.4, which completes the embedding of a single level.

As mentioned before, the theorem follows by incorporating the nesting techniques of Section 4.3. It is straightforward to verify the details. \square

6 Related notions of dimensionality

Theorem 6.1 (Euclidean embeddings). *The upper bounds for $\dim(G)$ in Theorems 5.8 and 4.10 hold also for $\dim_2(G)$.*

Proof. Consider a contractive, injective embedding φ of $G = (V, E)$ into \mathbb{Z}^d such that for some fixed $0 < \varepsilon < 1$, and every two distinct vertices u, v , their images $\varphi(u)$ and $\varphi(v)$ differ in at least εd coordinates. Our proof of Theorems 5.8 can be easily modified to yield such an embedding (for some universal constant ε) by applying appropriate Chernoff bounds when the coordinates are formed (see the application of Lemma 5.7 in Section 5.3, for instance). A similar modification to the proof of Theorem 4.10 results with ε that depends only on the size s of the excluded minor, namely, $\varepsilon \geq \Omega(1/4^s)$, because a modified proof of Theorem 4.2 guarantees that the images of every two distinct vertices are different in $\Omega(\rho)$, out of the $O(m^2\rho)$, coordinates.

After scaling φ by $(\varepsilon d)^{-\frac{1}{2}}$, this embedding satisfies:

1. $\|\varphi(u) - \varphi(v)\|_2 \geq 1$ for all $u \neq v \in V$; and
2. $\|\varphi(u) - \varphi(v)\|_2 \leq 1/\sqrt{\varepsilon}$ for all $(u, v) \in E$.

We now show that, for every fixed $0 < \nu < 1$, the constant $1/\sqrt{\varepsilon}$ can be reduced to $(1 + \nu)^3$ for an arbitrarily small constant $\nu > 0$, while increasing the dimension d only to $O(d)$, where the hidden constant depends only on ε and ν . Let $K = K(\varepsilon, \nu) \geq 1/\sqrt{\varepsilon}$ be determined later. For $x \in \mathbb{R}^d$, let $B_2(x, r) = \{y \in \mathbb{R}^d : \|x - y\|_2 < r\}$. Since, for every distinct $u, v \in V$, the balls $B_2(\varphi(u), \frac{1}{2})$ and $B_2(\varphi(v), \frac{1}{2})$ are disjoint, simple volume arguments show that, for all $u \in V$, $|\{v \in V : \|\varphi(v) - \varphi(u)\|_2 \leq 2K\}| \leq (c_1 K)^d$, where $c_1 > 0$ is a universal constant (constant independent of ε and ν).

We claim that there exists a map $\beta : V \rightarrow \mathbb{R}^{c_2 d}$, where $c_2 = c_2(K, \nu)$, satisfying

$$(*) \quad 1 \leq \|\beta(u) - \beta(v)\|_2 \leq 1 + \nu \quad \text{for } u \neq v \in V \text{ with } \|\varphi(u) - \varphi(v)\|_2 \leq K,$$

Assuming the existence of such a map β , we arrive at our final map $\gamma : V \rightarrow \mathbb{R}^{O(d)}$ defined by the direct sum $\gamma = \frac{1}{K}\varphi \oplus \beta$, which satisfies

- (1') $\|\gamma(u) - \gamma(v)\|_2 \geq 1$ for all $u \neq v \in V$; and
- (2') $\|\gamma(u) - \gamma(v)\|_2 \leq \sqrt{1/K^2\varepsilon + (1 + \nu)^2}$ for all $(u, v) \in E$.

Choosing $K(\varepsilon, \nu)$ to be sufficiently large, say $K = 1/\sqrt{\varepsilon\nu}$, we see that the bound in (2') is at most $(1 + \nu)^3$.

It remains now to prove the existence of such β . Let $c_2 = c_2(K, \nu)$ be determined later. We define $\beta(u)$ to be a random vector in $\mathbb{R}^{c_2 d}$, where each coordinate is chosen to be 0 or 1 uniformly at

random. We argue that if c_2 is large enough, then with positive probability, the required condition (*) holds, up to scaling, and this will complete the proof.

Let Y_{uv} be the number of coordinates in which $\beta(u)$ and $\beta(v)$ disagree. Clearly $\mathbb{E}[Y_{uv}] = \frac{1}{2}c_2d$. Furthermore, by standard Chernoff bounds, there exists $c_3 = c_3(\nu) > 0$ such that

$$\Pr \left[Y_{uv} < \frac{1}{1+\nu} \mathbb{E}[Y_{uv}] \text{ or } Y_{uv} > (1+\nu) \mathbb{E}[Y_{uv}] \right] \leq e^{-c_3c_2d}.$$

Let \mathcal{E}_u be the event that there exists some $\varphi(v) \in B_2(\varphi(u), K)$, for which either $Y_{uv} < \frac{1}{1+\nu} \mathbb{E}[Y_{uv}]$ or $Y_{uv} > (1+\nu) \mathbb{E}[Y_{uv}]$. Since $|\{v \in V : \|\varphi(v) - \varphi(u)\|_2 \leq K\}| \leq (c_1K)^d$, we can choose c_2 to be large enough so that (by a union bound) $\Pr[\mathcal{E}_u] \leq e^{-c_3c_2d/2}$. Finally, note that each event \mathcal{E}_u is mutually independent of all events \mathcal{E}_v for which $\varphi(v) \notin B_2(\varphi(u), 2K)$. The number of such events is again at most $(c_1K)^d$, and hence choosing $c_2 > 0$ to be large enough and applying the local lemma (Lemma 5.4), we see that, with positive probability, no event \mathcal{E}_u occurs. In this case, the map β satisfies $\sqrt{\frac{1}{1+\nu}c_2d/2} \leq \|\beta(u) - \beta(v)\|_2 \leq \sqrt{(1+\nu)c_2d/2}$ whenever $u \neq v$ and $\|\varphi(u) - \varphi(v)\|_2 \leq K$. Scaling this map proves the existence of the required β and completes the proof of the theorem. \square

Theorem 6.2. *If $G = (V, E)$ is a $\Theta(k)$ -degree expander with $1 \leq k \leq \log |V|$, then $\dim_2(G) = \Omega(\frac{\log |V|}{\log \log_k \log |V|})$. For a $\Theta(\log |V|)$ -degree expander, $\dim_2(G) = \Omega(\rho_G \log \rho_G)$.*

Proof. Similar to that of Theorem 3.2. \square

Acknowledgments

We are indebted to Nati Linial for his part in initiating this research. We are particularly grateful to him for valuable discussions in an early stage of the work and for his suggestion to focus on trees.

References

- [1] N. Alon and J. H. Spencer. *The probabilistic method*. Wiley-Interscience [John Wiley & Sons], New York, second edition, 2000.
- [2] P. Assouad. Plongements lipschitziens dans \mathbf{R}^n . *Bull. Soc. Math. France*, 111(4):429–448, 1983.
- [3] Y. Bartal. Probabilistic approximation of metric spaces and its algorithmic applications. In *37th Annual Symposium on Foundations of Computer Science*, pages 184–193. IEEE, 1996.
- [4] J. A. Bondy and U. S. R. Murty. *Graph theory with applications*. American Elsevier Publishing Co., Inc., New York, 1976.
- [5] A. Brandstädt, V. Chepoi, and F. Dragan. Distance approximating trees for chordal and dually chordal graphs. *J. Algorithms*, 30(1):166–184, 1999.
- [6] V. Chepoi and F. Dragan. A note on distance approximating trees in graphs. *European J. Combin.*, 21(6):761–766, 2000.
- [7] F. R. K. Chung. Labelings of graphs. In *Selected topics in graph theory*, 3, pages 151–168. Academic Press, San Diego, CA, 1988.
- [8] P. Erdős, F. Harary, and W. T. Tutte. On the dimension of a graph. *Mathematika*, 12: 118–122, 1965.

- [9] U. Feige. Approximating the bandwidth via volume respecting embeddings. *J. Comput. System Sci.*, 60(3): 510–539, 2000.
- [10] J. Fakcharoenphol and K. Talwar. An improved decomposition theorem for graphs excluding a fixed minor. *Proceedings of 6th Workshop on Approximation, Randomization, and Combinatorial Optimization*, Springer Lecture Notes in Computer Science 2764: 36–46, 2003.
- [11] A. Gupta, R. Krauthgamer, and J. R. Lee. Bounded geometries, fractals, and low-distortion embeddings. *Proceedings of the 44th Annual Symposium on Foundations of Computer Science*, 2003.
- [12] J. Heinonen. *Lectures on analysis on metric spaces*. Universitext. Springer-Verlag, New York, 2001.
- [13] P. Indyk. Algorithmic applications of low-distortion geometric embeddings. In *Proceedings of the 42nd Annual IEEE Symposium on Foundations of Computer Science*, pages 10–33. October 2001.
- [14] P. Klein, S. A. Plotkin, and S. Rao. Excluded minors, network decomposition, and multicommodity flow. In *25th Annual ACM Symposium on Theory of Computing*, pages 682–690, May 1993.
- [15] N. Linial. Variation on a theme of Levin. In J. Matoušek, Ed., *Open Problems, Workshop on Discrete Metric Spaces and their Algorithmic Applications*. Haifa, March 2002.
- [16] N. Linial, E. London, and Y. Rabinovich. The geometry of graphs and some of its algorithmic applications. *Combinatorica*, 15(2):215–245, 1995.
- [17] N. Linial and M. Saks. Low diameter graph decompositions. *Combinatorica*, 13(4):441–454, 1993.
- [18] L. Lovász and K. Vesztegombi. Geometric representations of graphs. In *Paul Erdős, Proc. Conf.*, Budapest, 1999.
- [19] J. Matoušek. *Lectures on discrete geometry*, Vol. 212 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2002.
- [20] R. Motwani and P. Raghavan. *Randomized Algorithms*. Cambridge University Press, 1995.
- [21] S. Rao. Small distortion and volume preserving embeddings for planar and Euclidean metrics. In *Proceedings of the 15th Annual Symposium on Computational Geometry*, pages 300–306. ACM, 1999.